

# iMedix: Your Personal Health Advisor.

## Incruse Ellipta Blister, With Inhalation Device

---

Search for medical details concerning Incruse Ellipta inhalation on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

**Side Effects** Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: difficult/painful urination. Rarely, this medication may cause sudden severe wheezing/trouble breathing right after use. If this occurs, use your quick-relief inhaler and get medical help right away. Get medical help right away if you have any very serious side effects, including: eye pain/swelling/redness, vision changes (such as seeing rainbows around lights at night, blurred vision). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Umeclidinium is used to control and prevent symptoms (such as wheezing, shortness of breath) caused by ongoing lung disease (chronic obstructive pulmonary disease-COPD, which includes bronchitis and emphysema). It works by relaxing the muscles around the airways so that they open up and you can breathe more easily.

Umeclidinium belongs to a class of drugs known as anticholinergics. Controlling symptoms of breathing problems can decrease time lost from work or school. This medication must be used regularly to be effective. It does not work right away and should not be used to relieve sudden shortness of breath. If wheezing or sudden shortness of breath occurs, use your quick-relief inhaler (such as albuterol, also called salbutamol in some countries) as prescribed. How to use Incruse Ellipta Blister, With Inhalation Device Read the Patient Information Leaflet and Instructions for Use Leaflet for directions on the proper use of this medication. If you have any questions, ask your doctor or pharmacist. Inhale this medication by mouth as directed by your doctor, usually once daily. Do not open the cover of the inhaler until you are ready to use it. Each time you open the cover, you prepare a dose of the medication. If you open and close the cover without inhaling the medication, you will lose the dose. To avoid wasting a dose, do not close the cover until after you have inhaled the medication. You may or may not taste/feel the drug when you inhale. Do not exhale into the device. Use this medication regularly to get the most benefit from it. To help you remember, use it at the same time every day. Do not use it more than one time each day. Do not increase your dose, use this

medication more often, or stop using it without first consulting your doctor. Your condition will not improve any faster, and your risk of side effects will increase. If you are using other inhalers at the same time, wait at least 1 minute between the use of each medication. Learn which of your inhalers you should use every day and which you should use if your breathing suddenly worsens (quick-relief drugs). Ask your doctor ahead of time what you should do if you have new or worsening cough or shortness of breath, wheezing, increased sputum, waking up at night with trouble breathing, if you use your quick-relief inhaler more often, or if your quick-relief inhaler does not seem to be working well. Learn when you can treat sudden breathing problems by yourself and when you must get medical help right away. Tell your doctor if your condition does not improve or if it worsens. Precautions Before using umeclidinium, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients (such as milk proteins), which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: personal or family history of glaucoma (angle-closure type), difficulty urinating (such as due to an enlarged prostate). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Avoid smoking and other things that can worsen breathing problems. Because the flu virus can also worsen breathing problems, ask your doctor or pharmacist if you should have a flu shot every year. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Do not use more than one dose each day. Store at room temperature away from light, heat, and moisture. Do not store in the bathroom. Do not open the sealed tray that the inhaler comes in until you are ready to use the new inhaler. Throw away the inhaler 6 weeks after first removing it from the tray or when the dose counter reads zero, whichever comes first. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

## **Side Effects**

---

## **Uses**

---

---

---

---

---

## Interactions

---

---

---

---

## Other Details

---

---