

# iMedix: Your Personal Health Advisor.

## Inderal LA

---

Search for medical details concerning Inderal LA oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

## Description

**Side Effects** Dizziness, lightheadedness, or tiredness may occur as your body adjusts to the medication. Nausea/vomiting, stomach pain, vision changes, trouble sleeping, and unusual dreams may also occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. To reduce the risk of dizziness and lightheadedness, get up slowly when rising from a sitting or lying position. This drug may reduce blood flow to your hands and feet, causing them to feel cold. Smoking may worsen this effect. Dress warmly and avoid tobacco use. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: blue fingers/toes, mental/mood changes (such as depression), numbness/tingling of arms/legs, new or worsening symptoms of heart failure (such as shortness of breath, swelling ankles/feet, unusual tiredness, unusual/sudden weight gain), very slow heartbeat, fainting, decreased sexual ability, increased thirst/urination, easy bruising/bleeding, signs of infection (such as sore throat that doesn't go away, fever), aching/swollen joints. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** This medication is a beta blocker used to treat high blood pressure, irregular heartbeats, shaking (tremors), and other conditions as determined by your doctor. It is used after a heart attack to improve survival. It is also used to prevent migraine headaches and chest pain (angina). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. Preventing chest pain can help improve your ability to exercise. This drug works by blocking the action of certain natural chemicals in your body (such as epinephrine) that affect the heart and blood vessels. This effect reduces heart rate, blood pressure, and strain on the heart. **How to use Inderal LA** Take this medication by mouth as directed by your doctor, usually once daily. Swallow the capsules whole; do not crush or chew the capsules. The dosage is based on your medical condition and response to treatment. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day. Keep taking this medication even if you feel well. This medication is used to help prevent chest pain or migraines. It should

not be used to treat chest pain or migraines when they occur. Use other medications (such as nitroglycerin tablets placed under the tongue for chest pain, sumatriptan for migraines) to relieve sudden attacks as directed by your doctor. Consult your doctor or pharmacist for details. It may take 1 to 2 weeks before you get the full benefit of this drug. If you also take certain drugs to lower your cholesterol (bile acid-binding resins such as cholestyramine or colestipol), take propranolol at least 1 hour before or at least 4 hours after these medications. Tell your doctor if your condition worsens (such as your routine blood pressure readings increase, your chest pain or migraines occur more often). Warnings Do not stop using this drug without first consulting your doctor. Your condition may become worse when the drug is suddenly stopped, especially if you have chest pain (angina) or heart disease (such as coronary artery disease, ischemic heart disease, high blood pressure). If your doctor decides you should no longer use this drug, you must gradually decrease your dose according to your doctor's instructions. When gradually stopping this medication, it is recommended that you temporarily limit physical activity to decrease strain on the heart. Get medical help right away if you develop: worsening chest pain, tightness/pressure in the chest, chest pain spreading to the jaw/neck/arm, unusual sweating, trouble breathing, or fast/irregular heartbeat. Precautions Before taking propranolol, tell your doctor or pharmacist if you are allergic to it; or if you have had a serious reaction to other beta blockers (such as metoprolol); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: other breathing problems (such as asthma, bronchitis, emphysema), heart failure, certain types of heart rhythm problems (sinus bradycardia, Wolff-Parkinson-White syndrome, second- or third-degree atrioventricular block), overactive thyroid (hyperthyroidism), kidney disease, liver disease, blood circulation problems (such as Raynaud's disease), a certain type of tumor (pheochromocytoma), mental/mood disorders (such as depression), certain muscle/nerve disease (myasthenia gravis), severe allergic reactions. This drug may make you dizzy. Alcohol or marijuana (cannabis) can make you more dizzy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). This product may prevent the fast/pounding heartbeat you would usually feel when your blood sugar level falls too low (hypoglycemia). The risk is higher if you have diabetes, or are vomiting, fasting, or not eating regularly. Other symptoms of low blood sugar level, such as dizziness and sweating, are not affected by this drug. If you have diabetes, this product may make it harder to control your blood sugar levels. Check your blood sugar levels regularly as directed by your doctor. Tell your doctor right away if you have symptoms of high blood sugar such as increased thirst/urination. Your doctor may need to adjust your diabetes medication, exercise program, or diet. Children may be at greater risk for low blood sugar (hypoglycemia) while using this drug, especially if they are vomiting or not eating regularly. To help prevent low blood sugar, make sure children eat regularly. If your child cannot eat regularly, is vomiting, or has symptoms of low blood sugar (such as sweating, shaking), tell the doctor right away. During pregnancy, this medication should be used only when clearly needed. Infants exposed to this medication during pregnancy may have low birth weight, low blood sugar, or slow breathing/heartbeat. Discuss the risks and benefits with your doctor. This drug passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include unusually slow heartbeat, severe dizziness, slow or shallow breathing, weakness, or fainting. Do not share this medication with others. Lifestyle changes such as stress reduction programs, exercise and dietary changes may increase the effectiveness of this medicine. Talk to your doctor or pharmacist about lifestyle changes that might benefit you. Have your blood pressure and pulse checked regularly while taking this medication. It may be best to learn how to monitor your own blood pressure and pulse. Discuss this with your doctor. If you miss a dose, take it as soon as you remember, but not if it is within 8 hours of the next dose. If it is within 8 hours of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal

company. **Interactions** Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: alpha blockers (such as prazosin), aluminum hydroxide, chlorpromazine, other drugs to treat high blood pressure (such as clonidine, hydralazine, methyldopa), epinephrine, fezolinetant, fingolimod, haloperidol, other heart medications (such as digoxin, disopyramide, propafenone, quinidine), mefloquine, rizatriptan, theophylline, thioridazine, thyroid hormones (such as levothyroxine), warfarin. Other medications can affect the removal of propranolol from your body, which may affect how propranolol works. Examples include cimetidine, St. John's wort, certain SSRI antidepressants (such as fluoxetine, fluvoxamine, paroxetine), rifamycins (such as rifabutin), among others. Some products have ingredients that could raise your heart rate or blood pressure. Tell your pharmacist what products you are using, and ask how to use them safely (especially cough-and-cold products, diet aids, or NSAIDs such as ibuprofen/naproxen). This medication may interfere with certain lab tests (such as glaucoma screening test), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug.

---

---

## **Side Effects**

---

---

## **Uses**

---

---

---

---

---

---

## **Interactions**

---

---

## **Other Details**

---

---