

# iMedix: Your Personal Health Advisor.

## Inveltys Suspension, Drops (Final Dose Form)

---

Search for medical details concerning Inveltys ophthalmic (eye) on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **Active Ingredient:**
- **Dosage Form:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

Side Effects Eye discomfort, redness, or temporary blurred vision may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: vision changes (such as decreased vision), eye pain. This medication may mask the signs of an eye infection. It may also increase the risk of developing an eye infection, especially with longer use. Tell your doctor right away if you have new or worsening symptoms such as eye discharge/swelling/redness. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used after eye surgery to help relieve pain and swelling (inflammation). Loteprednol belongs to a class of drugs known as corticosteroids. How to use Inveltys Suspension, Drops (Final Dose Form) Read the manufacturer's Instructions for Use if available from your pharmacist before you start using loteprednol and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Use this medication in the affected eye(s) as directed by your doctor, usually twice daily. To apply eye drops, wash hands first. Before using, shake the bottle well for 1 to 2 seconds. To avoid contamination, do not touch the dropper tip or let it touch your eye or any other surface. If you are wearing contact lenses, remove them before using eye drops. Wait at least 15 minutes before replacing your contact lenses. Tilt your head back, look upward, and pull down the lower eyelid to make a pouch. Hold the dropper directly over your eye and place one drop into the pouch. Look downward, gently close your eyes, and place one finger at the corner of your eye (near the nose). Apply gentle pressure for 1 to 2 minutes before opening your eyes. This will prevent the medication from draining out. Try not to blink or rub your eye. Repeat these steps if your dose is for more than one drop. If directed to use this medication in both eyes, repeat these steps for your other eye. Do not rinse the dropper. Replace the dropper cap after each use. If you are using another kind of eye medication (such as drops or ointments), wait at least 5 to 10 minutes before applying other medications. Use eye drops before eye ointments to allow the

drops to enter the eye. Use this medication regularly to get the most benefit from it. To help you remember, use it at the same times each day. The dosage and length of treatment are based on your medical condition and response to treatment. Continue to use this medication for the full time prescribed. Do not use this medication more often or for longer than prescribed. Doing so may increase your risk of side effects. Do not use this product if it becomes contaminated (for example, drops turn a dark color). Use of contaminated eye medication can cause infection, serious damage to the eye, and loss of vision. Tell your doctor if your condition does not get better or if it gets worse. Precautions Before using loteprednol, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: eye infections, cataracts, glaucoma. After you apply this drug, your vision may become temporarily blurred. Do not drive, use machinery, or do any activity that requires clear vision until you can do it safely. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for another eye condition unless told to do so by your doctor. A different medication may be necessary in that case. Lab and/or medical tests (such as eye exams) may be done while you are using this medication. Keep all medical and lab appointments. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not freeze. Do not store in the bathroom. Keep bottle upright and tightly closed when not in use. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

---

## **Side Effects**

---

---

---

## **Uses**

---

---

---

---

---

## Interactions

---

---

## Other Details

---

---