

iMedix: Your Personal Health Advisor.

Irbesartan-Hydrochlorothiazide

Search for medical details concerning irbesartan-hydrochlorothiazide oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Dizziness, lightheadedness, or upset stomach may occur as your body adjusts to the medication. If any of these effects last or get worse, tell your doctor or pharmacist promptly. To reduce the risk of dizziness and lightheadedness, get up slowly when rising from a sitting or lying position. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. This product may cause dehydration and electrolyte imbalance. Tell your doctor right away if you have any symptoms of dehydration or electrolyte imbalance, including: extreme thirst, very dry mouth, muscle cramps/weakness, fast/slow/irregular heartbeat, confusion. Tell your doctor right away if you have any serious side effects, including: fainting, decrease in vision, eye pain, symptoms of a high potassium blood level (such as muscle weakness, slow/irregular heartbeat), signs of kidney problems (such as change in the amount of urine). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345.

How to use Irbesartan-Hydrochlorothiazide Take this medication by mouth as directed by your doctor, usually once daily with or without food. If this medication causes you to urinate more often, it is best to take it at least 4 hours before your bedtime to prevent having to get up to urinate. If you also take certain drugs to lower your cholesterol (bile acid-binding resins such as cholestyramine or colestipol), take irbesartan/hydrochlorothiazide at least 4 hours before or at least 4 to 6 hours after these medications. The dosage is based on your medical condition and response to treatment. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day. Keep taking this medication even if you feel well. Most people with high blood pressure do not feel sick. Tell your doctor if your condition does not improve or if it worsens (for example, your blood pressure readings increase).

Precautions Before taking this medication, tell your doctor or pharmacist if you are allergic to irbesartan or hydrochlorothiazide; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: gout,

kidney disease, liver disease, lupus, dehydration, high level of potassium in the blood, skin cancer. This drug may make you dizzy. Alcohol or marijuana (cannabis) can make you more dizzy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Severe sweating, diarrhea, or vomiting can increase the risk for lightheadedness or dehydration. Report prolonged diarrhea or vomiting to your doctor. To prevent dehydration, drink plenty of fluids unless your doctor directs you otherwise. If you have diabetes, this product may affect your blood sugar. Check your blood sugar regularly as directed and share the results with your doctor. Your doctor may need to adjust your diabetes medication, exercise program, or diet. This medication may make you more sensitive to the sun. It may also increase your risk for skin cancer, especially if you take it for a long time. Limit your time in the sun. Avoid tanning booths and sunlamps. Use sunscreen and wear protective clothing when outdoors. Tell your doctor right away if you get sunburned, have skin blisters/redness, or notice new or changed moles/skin lesions. This product may affect your potassium levels. Before using potassium supplements or salt substitutes that contain potassium, consult your doctor or pharmacist. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Older adults may be more sensitive to the side effects of this drug, especially dizziness and change in the amount of urine (kidney problems). Tell your doctor if you are pregnant or plan to become pregnant. You should not become pregnant while using irbesartan/hydrochlorothiazide. This medication may harm an unborn baby. If you become pregnant, talk to your doctor right away about the risks and benefits of this medication. () It is unknown if irbesartan passes into breast milk. Hydrochlorothiazide passes into breast milk, but is unlikely to harm a nursing infant. Consult your doctor before breast-feeding. Consult your pharmacist or physician. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: aliskiren, dofetilide, lithium, drugs that may increase the level of potassium in the blood (such as sparsentan, ACE inhibitors including benazepril/lisinopril, birth control pills containing drospirenone). Some products have ingredients that could raise your blood pressure or worsen your heart failure. Tell your pharmacist what products you are using, and ask how to use them safely (especially cough-and-cold products, diet aids, or NSAIDs such as ibuprofen/naproxen). This medication may interfere with certain lab tests (such as parathyroid function), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: severe dizziness, fainting. Do not share this medication with others. Lifestyle changes such as stress reduction programs, exercise, and dietary changes may increase the effectiveness of this medicine. Talk to your doctor or pharmacist about lifestyle changes that might benefit you. Lab and/or medical tests (such as kidney function, potassium levels, skin exams) should be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details. Check your blood pressure regularly while taking this medication. Learn how to monitor your own blood pressure, and share the results with your doctor. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Side Effects

Uses

Interactions

Other Details
