

iMedix: Your Personal Health Advisor.

Isentress

Isentress, the brand name for raltegravir, is an antiretroviral medication used to treat HIV-1 in both adults and children. As an integrase inhibitor, it blocks the HIV integrase enzyme, preventing the viral DNA from integrating into the host's genome, crucial for viral replication.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Headache, nausea, and trouble sleeping may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. As your immune system gets stronger, it can begin to fight off infections you already had, possibly causing disease symptoms to come back. You could also have symptoms if your immune system becomes overactive. This reaction may happen at any time (soon after starting HIV treatment or many months later). Get medical help right away if you have any serious symptoms, including: unexplained weight loss, severe tiredness, muscle aches/weakness that doesn't go away, headaches that are severe or don't go away, joint pain, numbness/tingling of the hands/feet/arms/legs, vision changes, signs of infection (such as fever, chills, swollen lymph nodes, trouble breathing, cough, non-healing skin sores), signs of an overactive thyroid (such as irritability, nervousness, heat intolerance, fast/pounding/irregular heartbeat, bulging eyes, unusual growth in the neck/thyroid known as a goiter), signs of a certain nerve problem known as Guillain-Barre syndrome (such as unsteadiness, loss of coordination, trouble swallowing/speaking/chewing, trouble moving your eyes). Tell your doctor right away if you have any serious side effects, including: nausea/vomiting that doesn't stop, signs of kidney problems (such as change in the amount of urine), loss of appetite, severe stomach/abdominal pain, dark urine, yellowing eyes/skin. Changes in body fat may occur while you are taking this medication (such as increased fat in the upper back and stomach areas, decreased fat in the arms and legs). The cause and long-term effects of these changes are unknown. Discuss the risks and benefits of treatment with your doctor, as well as the possible use of exercise to reduce this side effect. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash (especially if you also have fever, blisters, mouth sores, eye redness/swelling, muscle/joint pain), itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Raltegravir is used with other HIV medications to help control HIV infection. It helps to decrease the amount of HIV in your body so your immune system can

work better. This lowers your chance of getting HIV complications (such as new infections, cancer) and improves your quality of life. Raltegravir is known as an integrase inhibitor. It blocks the virus from growing and infecting more cells. Raltegravir is not a cure for HIV infection. To decrease your risk of spreading HIV disease to others, continue to take all HIV medications exactly as prescribed by your doctor. Use an effective barrier method (latex or polyurethane condoms/dental dams) during sexual activity as directed by your doctor. Do not share personal items (such as needles/syringes, toothbrushes, and razors) that may have contacted blood or other body fluids. Consult your doctor or pharmacist for more details.

How to use Isentress oral

Read the Patient Information Leaflet if available from your pharmacist before you start taking raltegravir and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth with or without food as directed by your doctor, usually once or twice daily. The dosage is based on your medical condition, response to treatment, and other medications you may be taking. Be sure to tell your doctor and pharmacist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). The manufacturer directs to swallow this medication whole. However, many similar drugs (immediate-release tablets) can be split/crushed. Follow your doctor's directions on how to take this medication. Avoid using antacids that contain aluminum or magnesium while taking this medication. These antacids can make raltegravir work less well. It is very important to continue taking this medication (and other HIV medications) exactly as prescribed by your doctor. Do not skip any doses. Do not increase your dose or use this drug more often or for longer than prescribed. Your condition will not improve any faster, and your risk of side effects will increase. Do not take less of this drug than prescribed or stop taking it (or other HIV medicines) even for a short time unless directed to do so by your doctor. Doing so may cause the amount of virus to increase and/or make the infection more difficult to treat (resistant). For the best effect, take this medication at evenly spaced times. To help you remember, take this medication at the same time(s) every day. Do not switch between the film-coated tablet, the chewable tablet, or the powder packet for oral suspension forms of raltegravir without asking your doctor or pharmacist first.

Precautions

Before taking raltegravir, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver disease (such as hepatitis B, hepatitis C), muscle disorders (such as rhabdomyolysis, myopathy), high blood levels of creatine kinase (high CK test results). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Treatment can lower the risk of passing HIV infection to your baby, and raltegravir may be part of that treatment. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. Because breast milk can transmit HIV, do not breast-feed.

Interactions

Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. A product that may interact with this drug is: orlistat.

Overdose

If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Lab and/or medical tests (such as viral load, T-cell counts) should be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store in the original package at room temperature away from light and moisture. To help protect from moisture, keep the desiccant in the bottle. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Side Effects

Uses

Interactions

Other Details
