

# iMedix: Your Personal Health Advisor.

## Isradipine

---

Search for medical details concerning isradipine oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

## Description

**Side Effects** Dizziness, nausea, headache, tiredness, flushing, and swelling of the ankles/feet may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. To reduce dizziness and lightheadedness, get up slowly when rising from a sitting or lying position. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: fainting, fast/irregular/pounding heartbeat, vision changes. Some people who already have severe heart disease may rarely develop worsening chest pain or a heart attack after starting this medication or increasing the dose. Get medical help right away if you experience: worsening chest pain, symptoms of a heart attack (such as chest/jaw/left arm pain, shortness of breath, unusual sweating). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Isradipine is used with or without other medications to treat high blood pressure (hypertension). Lowering high blood pressure helps prevent strokes, heart attacks, and kidney problems. Isradipine is called a calcium channel blocker. It works by relaxing blood vessels so blood can flow more easily. How to use Isradipine Take this medication by mouth with or without food as directed by your doctor, usually twice daily. The dosage is based on your medical condition and response to treatment. Your doctor may gradually increase your dose. Follow your doctor's instructions carefully. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same times each day. Keep taking this medication even if you feel well. Most people with high blood pressure do not feel sick. It may take 2 to 4 weeks before you get the full benefit of this drug. Tell your doctor if your condition does not improve or if it worsens (for example, your blood pressure readings remain high or increase). Precautions Before taking isradipine, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: certain structural heart problem (aortic stenosis),

liver disease, kidney disease. This drug may make you dizzy. Alcohol or marijuana (cannabis) can make you more dizzy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. Consult your doctor before breast-feeding. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Other medications can affect the removal of isradipine from your body, which may affect how isradipine works. Examples include cimetidine, azole antifungals (such as itraconazole), macrolide antibiotics (such as amoxil, zithromax, flagyl, levofloxacin, ciprofloxacin), rifamycins (such as rifabutin), St. John's wort, drugs used to treat seizures (such as phenytoin), among others. Some products have ingredients that could raise your blood pressure. Tell your pharmacist what products you are using, and ask how to use them safely (especially cough-and-cold products, diet aids, or NSAIDs such as ibuprofen/naproxen). Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Talk with your doctor about making changes to your lifestyle that may help this medication work better (such as stress reduction programs, exercise, and dietary changes). Lab and/or medical tests (such as liver function, blood pressure, electrocardiograms) may be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details. There are different types of this medication available. They may not have the same effects. Do not change types without consulting your doctor or pharmacist. Have your blood pressure checked regularly while taking this medication. Discuss with your doctor how to monitor your own blood pressure. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

---

## **Side Effects**

---

---

---

## **Uses**

---

---

---

---

---

## Interactions

---

---

---

## Other Details

---

---