

# iMedix: Your Personal Health Advisor.

## Jaypirca 100 Mg Tablet Antineoplastic – Protein-Tyrosine Kinase Inhibitors

---

Search for medical details concerning Jaypirca oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

### Description

Side Effects Diarrhea, dizziness, muscle/joint pain, tiredness, or swelling may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. People using this medication may have serious side effects. However, you have been prescribed this drug because your doctor has judged that the benefit to you is greater than the risk of side effects. Careful monitoring by your doctor may decrease your risk. This medication may raise your blood pressure. Check your blood pressure regularly and tell your doctor if the results are high. Tell your doctor right away if you have any serious side effects, including: easy bruising/bleeding, fast/irregular heartbeat. Get medical help right away if you have any very serious side effects, including: coughing up blood, black/bloody stools, vomit that looks like coffee grounds, pink/bloody urine, confusion, severe headache, fainting, shortness of breath. This medication may increase the risk of other cancers (especially skin cancer). Limit your time in the sun. Use sunscreen and wear protective clothing when outdoors. Consult your doctor for more details. This medication may lower your ability to fight infections. This may make you more likely to get a serious (rarely fatal) infection or make any infection you have worse. Tell your doctor right away if you have any signs of infection (such as sore throat that doesn't go away, fever, chills, cough). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used to treat a certain type of cancer (mantle cell lymphoma-MCL). Pirtobrutinib belongs to a class of drugs known as kinase inhibitors. It works by slowing or stopping the growth of cancer cells. How to use Jaypirca 100 Mg Tablet Antineoplastic – Protein-Tyrosine Kinase Inhibitors Read the Patient Information Leaflet if available from your pharmacist before you start taking pirtobrutinib and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth with or without food as directed by your doctor, usually once daily. Swallow the tablets whole with water. Do not split, crush, or chew the tablets. The dosage is based on your medical condition, response to treatment, and other medications you may be taking. Be sure to tell your doctor and pharmacist about all the products you use (including

prescription drugs, nonprescription drugs, and herbal products). Use this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day. Do not increase your dose or use this drug more often or for longer than prescribed. Your condition will not improve any faster, and your risk of serious side effects will increase. Precautions Before taking pirtobrutinib, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: bleeding/blood problems, heart problems (such as irregular heartbeat), high blood pressure, kidney problems. This drug may make you dizzy. Alcohol or marijuana (cannabis) can make you more dizzy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana(cannabis). Pirtobrutinib can make you more likely to get infections or may make current infections worse. Stay away from anyone who has an infection that may easily spread (such as chickenpox, COVID-19, measles, flu). Talk to your doctor if you have been exposed to an infection or for more details. Tell your health care professional that you are using pirtobrutinib before having any immunizations/vaccinations. Avoid contact with people who have recently received live vaccines (such as flu vaccine inhaled through the nose). To lower the chance of getting cut, bruised, or injured, use caution with sharp objects like razors and nail cutters, and avoid activities such as contact sports. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Your doctor may tell you to temporarily stop treatment with this medication before certain medical, surgical, or dental procedures. Tell your doctor if you are pregnant or plan to become pregnant. You should not become pregnant while using pirtobrutinib. Pirtobrutinib may harm an unborn baby. Your doctor should order a pregnancy test before you start this medication. Ask about reliable forms of birth control while using this medication and for at least 1 week after the last dose. If you become pregnant, talk to your doctor right away about the risks and benefits of this medication. It is unknown if this drug passes into breast milk. Because of the possible risk to the infant, breast-feeding while using this drug and for at least 1 week after the last dose is not recommended. Consult your doctor before breast-feeding.

**Interactions** Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Other medications can affect the removal of pirtobrutinib from your body, which may affect how pirtobrutinib works. Examples include rifamycins (such as rifampin, rifabutin), certain drugs used to treat seizures (such as carbamazepine, phenytoin), St. John's wort, among others.

**Overdose** If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Lab and/or medical tests (such as complete blood count, kidney function) should be done before you start taking this medication and while you are taking it. Keep all medical and lab appointments. Consult your doctor for more details. If you miss a dose, take it as soon as you remember if it has been less than 12 hours since the time of your dose. If it has been more than 12 hours, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

## Side Effects

---

## Uses

---

---

---

---

---

---

## Interactions

---

---

## Other Details

---

---