

iMedix: Your Personal Health Advisor.

Linzess

Search for medical details concerning Linzess oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Diarrhea, bloating, gas, or stomach/abdominal pain/discomfort may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Stop taking this medication and tell your doctor right away if you have any serious side effects, including: severe diarrhea. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** Linaclotide is used to treat certain types of bowel problems (irritable bowel syndrome with constipation, chronic idiopathic constipation, functional constipation). It works by increasing fluid in your intestines and helping speed up movement of food through the gut. Linaclotide may improve stool texture and lessen symptoms such as bloating, abdominal pain/discomfort, straining, and feelings of incomplete bowel movements. **How to use Linzess** Read the Medication Guide provided by your pharmacist before you start taking linaclotide and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth as directed by your doctor, usually once daily on an empty stomach, at least 30 minutes before a meal. Swallow the capsules whole. Do not break or chew the capsules. If you have trouble swallowing the capsule, it may be opened and the contents mixed in a teaspoonful of applesauce. Swallow the mixture right away, without chewing it. Do not save the mixture for future use. You may also empty the capsule's content into an ounce (30 milliliters) of water. Swirl the mixture for 20 seconds and drink right away, without chewing it. Add another ounce (30 milliliters) of water to the cup, swirl for 20 seconds and drink right away to get your whole dose. The water mixture may also be given through a tube into the stomach (nasogastric or gastric tube). If you are giving this medication through a tube, ask your health care professional for detailed instructions on how to give it. The dosage is based on your medical condition and response to treatment. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day. Tell your doctor if your condition does not improve or if it worsens. **Warnings** This medication must not be used by children younger than 2 years due to the risk for

harm (especially serious dehydration). Talk to your doctor for details. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. A product that may interact with this drug is: pramlintide. Precautions Before taking linaclotide, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: stomach/intestinal blockage. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. If you miss a dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store in the original container at room temperature away from light and moisture. To help protect from moisture, keep the desiccant in the tightly closed container. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Side Effects

Uses

Interactions

Other Details

