

# iMedix: Your Personal Health Advisor.

## Lotrimin Ultra Cream

---

Search for medical details concerning Lotrimin Ultra topical on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

**Side Effects** Burning, irritation, redness, or itching at the application site may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. If your doctor has prescribed this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: blistering/swelling/oozing at the application site. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** This medication is used to treat a variety of fungal skin infections such as ringworm, athlete's foot, and jock itch. This medication is also used to treat a skin condition known as pityriasis (tinea versicolor), a fungal infection that causes a lightening or darkening of the skin of the neck, chest, arms, or legs. Butenafine is an antifungal that works by preventing the growth of fungus. Do not use this medication for the treatment of fungal nail infections. **How to use Lotrimin Ultra Cream** Use this medication on the skin only. Clean and thoroughly dry the area to be treated. Apply a thin layer of the medication to and around the affected area and gently rub in, usually 1 to 2 times daily as directed on the product package. If you have any questions, ask your doctor or pharmacist. If your doctor has prescribed this medication to you, then use as directed. Wash your hands after using unless the area to be treated includes the hands. Do not wrap, cover, or bandage the area unless directed to do so by your doctor. Do not apply the medication in the eyes, nose, or mouth, inside the vagina, or on the scalp. If you do get the medication in those areas, flush with plenty of water. The dosage and length of treatment depends on the type of infection being treated. Do not apply more often or use longer than directed. This may increase the risk of side effects. Use this medication regularly to get the most benefit from it. To help you remember, use it at the same time(s) each day. Continue to use this medication until the full treatment period is finished, even if symptoms disappear after a few days. Stopping the medication too early may result in a return of the infection. Inform your doctor if your condition worsens or does not improve after the treatment period. **Precautions** Before using this medication, tell your doctor or pharmacist if you are allergic to butenafine; or to allylamine antifungals (such as naftifine, terbinafine); or if

you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: athlete's foot on the bottom/side of the foot. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. If your doctor has directed you to use this medication, use it for your current condition only. Do not use it later for another infection unless your doctor tells you to. If you are treating athlete's foot, make sure to wear well-fitting, ventilated shoes. Also, change shoes and socks at least once daily. When this medication is used to treat pityriasis, your normal skin color may not return right away. It may take several months after treatment is completed for your natural skin color to return. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

---

---

## **Side Effects**

---

---

## **Uses**

---

---

---

---

---

---

---

## **Interactions**

---

---

## **Other Details**

