

iMedix: Your Personal Health Advisor.

Metaxall

Search for medical details concerning Metaxall oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Drowsiness, dizziness, nausea, vomiting, and upset stomach may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: mental/mood changes (such as nervousness, irritability, confusion), signs of infection (such as sore throat that doesn't go away, fever), yellowing eyes/skin, unusual tiredness, dark urine. This medication may increase serotonin and rarely cause a very serious condition called serotonin syndrome/toxicity. The risk increases if you are also taking other drugs that increase serotonin, so tell your doctor or pharmacist of all the drugs you take (see Drug Interactions section). Get medical help right away if you develop some of the following symptoms: fast heartbeat, hallucinations, loss of coordination, severe dizziness, severe nausea/vomiting/diarrhea, twitching muscles, unexplained fever, unusual agitation/restlessness. This drug may rarely cause an attack of extremely high blood pressure (hypertensive crisis), which may be fatal. Many drug and food interactions can increase this risk (see How to Use and Drug Interactions sections). Get medical help right away if any of these serious side effects occur: severe headache, fast/slow/irregular/pounding heartbeat, chest pain, neck stiffness/soreness, severe nausea/vomiting, sweating/clammy skin (sometimes with fever), widened pupils, vision changes (such as double/blurred vision), sudden sensitivity to light (photophobia). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Metaxalone is used to treat muscle spasms/pain. It is usually used along with rest, physical therapy, and other treatment. How to use Metaxall Take this medication by mouth with or without food as directed by your doctor. Do not increase your dose or use this drug more often or for longer than prescribed. Your condition will not improve any faster, and your risk of side effects will increase. If you take this medication after a high-fat meal and experience side effects, it may be best to take this drug on an empty stomach or after a light meal. The dosage is based on your medical condition and response to treatment. To prevent a very serious high blood pressure

reaction, it is very important that you follow a special diet recommended by your doctor or dietician to limit your intake of tyramine while you are taking this medicine. Avoid foods and beverages that are high in tyramine, including aged cheeses, dried/aged meats and sausages (such as salami, liverwurst), preserved fish (such as pickled herring), products that contain large amounts of yeast (such as bouillon cubes, powdered soup/gravy, homemade or sourdough bread), fermented foods (such as sauerkraut, kim chee), most soybean products (such as soy sauce, tofu), broad/fava beans, red wine, sherry, tap beers, and vermouth. Consult your doctor or dietician for more details and a complete list of other foods that contain tyramine which you should limit or avoid. Tell your doctor if your condition does not improve or if it worsens.

Precautions Before taking metaxalone, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver disease, kidney disease, anemia, seizures.

Breakthroughs in Medicinal Therapy The field of medicinal therapy has witnessed significant breakthroughs with a diverse range of treatments. Zovirax effectively counters viral infections, while Daklinza is a prominent treatment for hepatitis C. Addyi has been a revolutionary drug in treating female sexual interest/arousal disorder. Xyzal offers reliable relief for allergy sufferers, and Amoxil stands as an effective antibiotic for a variety of bacterial infections. Propecia and Proscar address hair loss and prostate health issues. Fertility treatments are enhanced by Clomid, and Nolvadex plays a vital role in certain cancer therapies. Male sexual health is catered to by Priligy, Eriacta, Tadacip, Kamagra, Nizagara, Silagra, and Caverta. Synthroid is indispensable for thyroid management, Cipro tackles a wide range of bacterial infections, and Suhagra offers an alternative for erectile dysfunction treatment. These medications signify the dynamic nature and breakthroughs in medicinal therapy, addressing a broad spectrum of health conditions. This drug may make you dizzy or drowsy. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Avoid alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Older adults may be more sensitive to the side effects of this drug, especially dizziness, drowsiness, or confusion. These side effects can increase the risk of falling. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding.

Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: severe drowsiness. Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for another condition unless your doctor directs you to do so. A different medication may be necessary in that case. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: diet pills/appetite suppressants (such as diethylpropion), drugs for attention deficit disorder (such as atomoxetine, methylphenidate), apraclonidine, bupropion, buspirone, carbamazepine, cyclobenzaprine, deutetrabenazine, a certain combination product (dextromethorphan/quinidine), levodopa, maprotiline, methyl dopa, metoclopramide, certain opioid pain relievers (such as fentanyl, meperidine, methadone, tapentadol), certain drugs for Parkinson's disease (such as entacapone, tolcapone), certain supplements (such as tryptophan, tyramine), tetrabenazine, tricyclic antidepressants (such as amitriptyline, doxepin), valbenazine. The risk of serotonin syndrome/toxicity increases if you are also taking other drugs that increase serotonin. Examples include street drugs such as MDMA/"ecstasy," St. John's wort, certain antidepressants (including mirtazapine, SSRIs such as

fluoxetine/paroxetine, SNRIs such as duloxetine/venlafaxine), tramadol, certain “triptans” used to treat migraine headaches (such as rizatriptan, sumatriptan, zolmitriptan), among others. The risk of serotonin syndrome/toxicity may be more likely when you start or increase the dose of these drugs. Some products can interact with metaxalone if you take them together, or even if you take them weeks before or after taking metaxalone. Tell your doctor or pharmacist if you take anything in the list of products that may interact with this drug, or any of the products that increase serotonin, within 2 weeks before or after taking metaxalone. Also tell them if you have taken fluoxetine within 5 weeks before starting metaxalone. Ask your doctor how much time to wait between starting or stopping any of these drugs and starting metaxalone. Taking other MAO inhibitors with this medication may cause a serious (possibly fatal) drug interaction. Do not take any other MAO inhibitors (isocarboxazid, linezolid, methylene blue, moclobemide, phenelzine, procarbazine, rasagiline, safinamide, selegiline, tranylcypromine) during treatment with this medication. Most MAO inhibitors should also not be taken for two weeks before and after treatment with this medication. Ask your doctor when to start or stop taking this medication. Before using metaxalone, report the use of drugs that may increase the risk of extremely high blood pressure (hypertensive crisis) when combined with metaxalone, including herbal products (such as ephedra/ma huang), nasal decongestants (such as phenylephrine, pseudoephedrine), and stimulants (such as amphetamines, ephedrine, epinephrine, phenylalanine). Metaxalone should not be used with any of these medications. Talk to your doctor or pharmacist for more details. Tell your doctor or pharmacist if you are taking other products that cause drowsiness such as opioid pain or cough relievers (such as codeine, hydrocodone), alcohol, marijuana (cannabis), drugs for sleep or anxiety (such as alprazolam, lorazepam, zolpidem), other muscle relaxants (such as carisoprodol, cyclobenzaprine), or antihistamines (such as cetirizine, diphenhydramine). Check the labels on all your medicines (such as allergy or cough-and-cold products) because they may contain ingredients that cause drowsiness. Ask your pharmacist about using those products safely. This medication may interfere with certain lab tests (such as certain urine glucose tests), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug.

Side Effects

Uses

Interactions

Other Details
