

# iMedix: Your Personal Health Advisor.

## Milk Of Magnesia Suspension, (Final Dose Form)

---

Search for medical details concerning Milk of Magnesia oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **Active Ingredient:**
- **Dosage Form:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

Side Effects Diarrhea may occur. If this effect lasts or gets worse, tell your doctor or pharmacist promptly. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: symptoms of high magnesium levels (such as muscle weakness, slow/irregular heartbeat, slow/shallow breathing, mental/mood changes such as confusion), symptoms of dehydration (such as decreased urination, dizziness, extreme thirst, very dry mouth), stomach/abdominal pain, bloody stools, rectal bleeding. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used for a short time to treat occasional constipation. It is a laxative (osmotic-type) that is thought to work by drawing water into the intestines, an effect that helps to cause movement of the intestines. This medication is also used to treat symptoms caused by too much stomach acid such as heartburn, upset stomach, or indigestion. It is an antacid that works by lowering the amount of acid in the stomach. How to use Milk Of Magnesia Suspension, (Final Dose Form) Take this product by mouth as directed. For the chewable form, chew thoroughly before swallowing. For the liquid form, shake the bottle well before each dose. Carefully measure the dose using a special measuring device/spoon. Do not use a household spoon because you may not get the correct dose. If you are taking this medication for constipation, drink a full glass of water (8 ounces or 240 milliliters) with each dose. Follow all directions on the product package, or use as directed by your doctor. If you have any questions, ask your doctor or pharmacist. Dosage is based on your medical condition and response to treatment. Extended use or overuse of this medication for constipation may result in dependence on laxatives and ongoing constipation. Overuse may also cause diarrhea that doesn't stop, dehydration, and mineral imbalances (such as high magnesium). Tell your doctor if your condition lasts or gets worse. For constipation problems, it may take 30 minutes to 6 hours to produce a bowel movement. Tell your doctor if this product does not produce a bowel movement, if you need to use this medication or other laxative products regularly for more than 1 week, or if you have rectal bleeding. For

stomach acid problems, do not take the maximum dose of this medication for more than 2 weeks unless directed by your doctor. If you think you may have a serious medical problem, get medical help right away. Precautions Before taking magnesium hydroxide, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney disease, appendicitis or symptoms of appendicitis (such as stomach/abdominal pain, nausea/vomiting), magnesium-restricted diet, sudden change in bowel habits that lasts for longer than 2 weeks. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: diarrhea that is severe or doesn't stop, muscle weakness, mental/mood changes (such as confusion), slow/irregular heartbeat, decreased urination, dizziness. Keep all medical and lab appointments. Lifestyle changes such as regular exercise and diet changes (including drinking enough water, eating a proper diet with fiber-rich foods such as bran, fresh fruits/vegetables) may prevent or relieve constipation. Lifestyle changes such as stress reduction programs, stopping smoking, limiting alcohol, and diet changes (such as avoiding caffeine/certain spices) may help to reduce heartburn and other stomach acid problems. Talk to your doctor or pharmacist about lifestyle changes that might benefit you. Not applicable. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: raltegravir, sodium polystyrene sulfonate. Magnesium hydroxide can decrease the absorption of other drugs such as dasatinib, atazanavir, gabapentin, digoxin, mycophenolate, phosphate supplements (such as potassium phosphate), tetracycline antibiotics (such as amoxil, zithromax, flagyl, levofloxacin, doxycycline, minocycline), certain azole antifungals (ketoconazole, itraconazole), and quinolone antibiotics (such as ciprofloxacin, levofloxacin). Separate your doses of these medications as far as possible from your doses of magnesium hydroxide. Ask your doctor or pharmacist about how long you should wait between doses and for help finding a dosing schedule that will work with all your medications.

---

---

---

## Side Effects

---

---

---

## Uses

---

---

---

---

---

---

## Interactions

---

---

---

---

---

---

## Other Details

---

---

---