

iMedix: Your Personal Health Advisor.

Myrbetriq

Myrbetriq, with its active ingredient mirabegron, serves as a novel treatment for overactive bladder (OAB) by targeting beta-3 adrenergic receptors in the bladder muscle, leading to increased bladder capacity and reduced symptoms of urgency and frequency.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Nausea, dizziness, fast heartbeat, runny/stuffy nose, or headache may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. This medication may raise your blood pressure. Check your blood pressure regularly and tell your doctor if the results are high. Tell your doctor right away if you have any serious side effects, including: burning/painful urination. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used to treat certain bladder problems (overactive bladder, neurogenic detrusor overactivity). Overactive bladder is a problem with how your bladder stores urine. Neurogenic detrusor overactivity is a bladder control condition caused by brain, spinal cord, or nerve problems. Symptoms of these conditions may include frequent urination, strong sudden urges to urinate that are hard to control, or involuntary loss of urine (incontinence). Mirabegron works by relaxing a certain bladder muscle (detrusor), which helps the bladder hold more urine and lessens symptoms of overactive bladder and neurogenic detrusor overactivity. How to use Myrbetriq oral Read the Patient Information Leaflet if available from your pharmacist before you start taking mirabegron and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth as directed by your doctor, usually once daily. Adults may take this medication with or without food. Children should take this medication with food. Swallow this medication whole. Do not crush, chew, or divide the medication. Doing so can release all of the drug at once, increasing the risk of side effects. The dosage is based on your medical condition and response to treatment. Children's dose is also based on weight. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day. Tell your doctor if your condition does not get better or if it gets worse. Precautions Before taking mirabegron, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or

other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: other bladder problems (such as blockage), high blood pressure, kidney problems, liver problems. This drug may make you dizzy. Alcohol or marijuana (cannabis) can make you more dizzy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Lab and/or medical tests (such as blood pressure) may be done while you are taking this medication. Keep all medical and lab appointments. If you miss a dose and it is within 12 hours of the time you usually take the dose, take it as soon as you remember. If it is more than 12 hours from the usual time of your dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. This medication can slow down the removal of other medications from your body, which may affect how they work. Examples of affected drugs include flecainide, propafenone, thioridazine, among others.

Side Effects

Uses

Interactions

Other Details
