

# iMedix: Your Personal Health Advisor.

## Nateglinide

---

Search for medical details concerning nateglinide oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

## Description

**Side Effects** Weight gain may occur. If this effect lasts or gets worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Nateglinide can cause low blood sugar (hypoglycemia) especially if you are taking other medicines for diabetes. Consuming large quantities of alcohol, not getting enough calories from food, or doing unusually heavy exercise may also lead to low blood sugar. Symptoms may include chills, cold sweat, dizziness, drowsiness, shaking, fast heartbeat, weakness, headache, fainting, tingling of the hands or feet, or hunger. It is a good habit to carry glucose tablets or gel to treat low blood sugar. If you don't have these reliable forms of glucose, raise your blood sugar quickly by eating a quick source of sugar such as table sugar, honey, candy, or drinking a glass of fruit juice or non-diet soda. Check with your doctor or pharmacist to find out what you should do if you miss a meal. Symptoms of high blood sugar (hyperglycemia) include increased thirst/urination. If these symptoms occur, tell your doctor right away. Your doctor may need to adjust your diabetes medication, exercise program, or diet. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Nateglinide is used alone or with other medications to control high blood sugar along with a proper diet and exercise program. It is used in people with type 2 diabetes. Controlling high blood sugar helps prevent kidney damage, blindness, nerve problems, loss of limbs, and sexual function problems. Proper control of diabetes may also lessen your risk of a heart attack or stroke. It works by stimulating the body to produce more insulin. Insulin is a natural substance that allows the body to properly use sugar from the diet. How to use Nateglinide Read the Patient Information leaflet if one is available from your pharmacist before you start using nateglinide and each time you get a refill. If you have any questions, consult your doctor or pharmacist. Take this medication by mouth 1 to 30 minutes before each main meal as directed by your doctor, usually 3 times daily. Take this drug no earlier than 30 minutes before the meal. Do not take a dose of medication if you are skipping that meal. If you are having liquid meals, a higher dose of this medication may be necessary

to control your blood sugar. Consult your doctor for more details. The dosage is based on your medical condition, number of meals per day, and response to treatment. Use this medication regularly as directed by your doctor in order to get the most benefit from it. Carefully follow the medication treatment plan, meal plan, and exercise program your doctor has recommended. Check your blood sugar regularly as directed by your doctor. Keep track of the results, and share them with your doctor. This is very important in order to determine the correct dose. Tell your doctor if your blood sugar measurements are too high or too low. Your treatment plan may need to be changed.

**Precautions** Before taking nateglinide, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney disease, liver disease, gout. You may experience blurred vision, dizziness, or drowsiness due to extremely low or high blood sugar. Do not drive, use machinery, or do any activity that requires alertness or clear vision until you are sure you can perform such activities safely. Limit alcohol while taking this medication because it can increase the risk of developing low blood sugar. During times of stress, such as fever, infection, injury, or surgery, it may be more difficult to control your blood sugar. Consult your doctor because increased stress may require a change in your treatment plan, medications, or blood sugar testing. During pregnancy, this medication should be used only when clearly needed. Pregnancy may cause or worsen diabetes. Discuss a plan with your doctor for managing your blood sugar while pregnant. Your doctor may change your diabetes treatment during your pregnancy. Discuss the risks and benefits of different treatments (such as diet, exercise, and medications including insulin). It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding.

**Overdose** If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: very fast heartbeat, vision changes, unexplained heavy sweating, agitation, fainting, seizures. Do not share this medication with others. Attend a diabetes education program to learn more about how to manage your diabetes with medications, diet, exercise, and regular medical exams. Learn the symptoms of high and low blood sugar and how to treat low blood sugar. Check your blood sugar regularly as directed and share the results with your doctor. Lab and/or medical tests (such as kidney/liver function tests, fasting blood glucose, hemoglobin A1c) should be done while you are taking this medication. Keep all medical and lab appointments. If you miss a dose, skip that dose. Take your next dose with your next meal. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

**Interactions** Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Many drugs can affect your blood sugar, making it harder to control. Examples include corticosteroids (such as prednisone), psychiatric medicines (such as olanzapine), fluoroquinolone antibiotics (such as amoxicil, zithromax, flagyl, levofloxacin, ciprofloxacin), among others. Check your blood sugar regularly as directed and share the results with your doctor. Tell your doctor right away if you have symptoms of high or low blood sugar. ( ) Your doctor may need to adjust your diabetes medication, exercise program, or diet. Check the labels on all your nonprescription medicines (such as cough-and-cold products) because they may contain ingredients that could affect your blood sugar. Ask your pharmacist about the safe use of those products. Beta-blocker medications (such as metoprolol, propranolol, glaucoma eye drops such as timolol) may prevent the fast/pounding heartbeat you would usually feel when your blood sugar falls too low (hypoglycemia). Other symptoms of low blood sugar such as dizziness, hunger, or sweating are unaffected by these drugs. Other medications can affect the results of urine tests for sugar or ketones. Consult your doctor or pharmacist for more information.

---

---

## **Side Effects**

---

## **Uses**

---

---

---

---

---

---

---

---

## **Interactions**

---

---

## **Other Details**

---

---