

iMedix: Your Personal Health Advisor.

Neo-Polycin HC Ointment

Search for medical details concerning Neo-Polycin HC ophthalmic (eye) on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Stinging/burning of the eyes for a minute or two or temporary blurred vision may occur . If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Use of this medication for prolonged/repeated periods may result in a new fungal eye infection and may increase your risk for other eye problems (such as glaucoma, cataracts). Do not use this medication for longer than prescribed. Tell your doctor right away if you notice any of the following: new or worsening eye symptoms (such as discharge, swelling, redness), vision problems, eye pain. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used to treat or prevent bacterial eye infections. This product contains neomycin, bacitracin, and polymyxin, which are antibiotics that work by stopping the growth of bacteria. It also contains hydrocortisone, which is an anti-inflammatory corticosteroid that works by reducing swelling. This medication treats/prevents only bacterial eye infections. It will not work for other types of eye infections. Unnecessary use or overuse of any antibiotic can lead to its decreased effectiveness. How to use Neo-Polycin HC Ointment Apply this medication to the eye(s) as directed by your doctor, usually every 3 or 4 hours. Do not wear contact lenses while you are using this medicine. Sterilize contact lenses according to manufacturer's directions, and check with your doctor before you begin using them again. To apply the eye ointment, wash your hands first. To avoid contamination, be careful not to touch the tube tip or let it touch your eye. Tilt your head back, look upward, and pull down the lower eyelid to make a pouch. Place a 1/2 inch (1.5 centimeters) strip of ointment into the pouch by squeezing the tube gently. Close the eye and roll the eyeball in all directions to spread the medication. Try not to blink and do not rub the eye. Repeat these steps for your other eye if so directed. Wipe the tip of the ointment tube with a clean tissue to remove excess medication before recapping it. Apply as often as directed by your doctor. If you are using another kind of eye medication (such as drops or ointments), apply the eye drops first and wait at least 5 minutes before applying this eye ointment. Use eye

drops before eye ointments to allow the eye drops to enter the eye. If you are using another type of eye ointment, wait at least 10 minutes after using this medication before applying it. Use this medication regularly in order to get the most benefit from it. Remember to use it at the same times each day. Continue to use this medication for the full time prescribed, even if symptoms disappear after a few days. Stopping the medication too early may allow bacteria to continue to grow, which may result in a relapse of the infection. Inform your doctor if there is no improvement in your condition after 2 days of treatment with this medication. **Precautions** Before using this medication, tell your doctor or pharmacist if you are allergic to neomycin, bacitracin, polymyxin, or hydrocortisone; or to aminoglycoside antibiotics (such as gentamicin, tobramycin); or to corticosteroids (such as dexamethasone, prednisone); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: other eye problems (such as glaucoma). After you apply this drug, your vision may become temporarily blurred or unstable. Do not drive, use machinery, or do any activity that requires clear vision until you can do it safely. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. **Overdose** This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. This medication has been prescribed for your current condition only. Throw away the unused medication after treatment is completed. Do not use it later for another infection unless your doctor tells you to. If you are using this medication for 10 days or more, lab and/or medical tests (such as eye exams) should be done while you are using this medication. Keep all medical and lab appointments. Consult your doctor for more details. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from heat. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. **Interactions** Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Side Effects

Uses

Interactions

Other Details
