iMedix: Your Personal Health Advisor.

Nexium

Search for medical details concerning Nexium oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings. Nexium 24HR is a medication used to treat frequent heartburn, a symptom of acid reflux. It works by reducing the amount of acid produced in the stomach, providing 24-hour relief from heartburn symptoms. Nexium 24HR comes in a delayed-release capsule and is available over-the-counter for convenient access.

- ActiveIngredient:
- DosageForm:
- Dosage:
- Indications:
- Manufacturer:
- Storage:
- Market Price:
- Drug Status:

Description

Side Effects Headache or abdominal pain may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: symptoms of a low magnesium blood level (such as muscle spasms, irregular heartbeat, seizures), signs of lupus (such as rash on nose and cheeks, new or worsening joint pain). This medication may rarely cause a severe intestinal condition due to a bacteria called C. difficile. This condition may occur during treatment or weeks to months after treatment has stopped. Tell your doctor right away if you develop: diarrhea that doesn't stop, abdominal or stomach pain/cramping, blood/mucus in your stool. If you have these symptoms, do not use anti-diarrhea or opioid products because they may make symptoms worse. Rarely, proton pump inhibitors (such as esomeprazole) have caused vitamin B-12 deficiency. The risk is increased if they are taken every day for a long time (3 years or longer). Tell your doctor right away if you develop symptoms of vitamin B-12 deficiency (such as unusual weakness, sore tongue, or numbness/tingling of the hands/feet). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: fever, swollen lymph nodes, rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing, signs of kidney problems (such as change in the amount of urine). Uses Esomeprazole is used to treat certain stomach and esophagus problems (such as acid reflux, ulcers). It works by decreasing the amount of acid your stomach makes. It relieves symptoms such as heartburn, difficulty swallowing, and cough. This medication helps heal acid damage to the stomach and esophagus, helps prevent ulcers, and may help prevent cancer of the esophagus. Esomeprazole belongs to a class of drugs known as proton pump inhibitors (PPIs). If you are self-treating with this medication, over-thecounter esomeprazole products are used to treat frequent heartburn (occurring 2 or more days a week). Since it may take 1 to 4 days to have full effect, these products do not relieve heartburn right away. For over-thecounter products, carefully read the package instructions to make sure the product is right for you. Check the ingredients on the label even if you have used the product before. The manufacturer may have changed the ingredients. Also, products with similar brand names may contain different ingredients meant for different

purposes. Taking the wrong product could harm you. How to use Nexium Capsule, Delayed Release (Enteric Coated) Read the Medication Guide and the Patient Information Leaflet if available from your pharmacist before you start taking esomeprazole and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth as directed by your doctor, usually once daily, at least 1 hour before a meal. If you are self-treating, follow all directions on the product package. The dosage and length of treatment are based on your medical condition and response to treatment. Do not increase your dose or take this drug more often than directed. If you have any questions, ask your doctor or pharmacist. Do not crush or chew this medication. Swallow the capsules whole. If you have difficulty swallowing this medication whole, you may open the capsule and sprinkle the contents into a tablespoon of unheated applesauce. Swallow the applesauce mixture right away without chewing it. Do not prepare the mixture ahead of time for later use. Doing so may destroy the drug. If needed, antacids may be taken along with this medication. If you are also taking sucralfate, take esomeprazole at least 30 minutes before sucralfate. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day. Continue to take this medication for the prescribed length of treatment even if you are feeling better. If you are self-treating with the over-the-counter product, do not take it for more than 14 days unless directed by your doctor. Tell your doctor if your condition lasts or gets worse. If you are self-treating, tell your doctor if your heartburn lasts after 14 days or if you need to use this medication more than once every 4 months. The risk of side effects goes up over time. Ask your doctor how long you should take this medication. If you think you may have a serious medical problem, get medical help right away. How long does it take for this medicine to take effect? The onset of action for Nexium is not immediate. It may take several days of consistent use to experience full relief of symptoms. Some patients may start to notice symptom improvement within the first 24 hours of taking the medication, but maximum effects typically occur after continuous use for 1 to 4 days. How long do the effects of this medicine last? The effects of Nexium can last for up to 24 hours, making it suitable for once-daily dosing. The duration of effect is due to the medication's ability to irreversibly inhibit the proton pump in stomach cells, which is responsible for acid secretion. Is it safe to consume alcohol while taking this medicine? While there are no direct contraindications against consuming alcohol while taking Nexium, alcohol can irritate the stomach lining and exacerbate symptoms of GERD and acid reflux. Patients are often advised to limit or avoid alcohol consumption to manage their symptoms more effectively. Is this a habit forming medicine? Nexium is not known to be habit-forming or to cause dependency. However, long-term use of proton pump inhibitors, including Nexium, should be managed and monitored by a healthcare provider due to potential risks associated with extended use, such as nutrient malabsorption or increased risk of certain infections. Can this medicine be taken during pregnancy? Nexium is classified as a pregnancy category B drug by the FDA, indicating that there is no evidence of harm to the fetus in animal studies, but well-controlled studies in pregnant women are lacking. Nexium should only be used during pregnancy if clearly needed and prescribed by a healthcare provider. Can this medicine be taken while breast-feeding? Esomeprazole is excreted into breast milk in small amounts. While there have been no reported adverse effects on nursing infants, caution is advised. The benefits of using Nexium while breast-feeding should be weighed against the potential risks to the infant, and use should be discussed with a healthcare provider. Precautions Before taking esomeprazole, tell your doctor or pharmacist if you are allergic to it; or to similar drugs (such as lansoprazole, omeprazole); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver disease, lupus. Some symptoms may actually be signs of a more serious condition. Get medical help right away if you have: heartburn with lightheadedness/sweating/dizziness, chest/jaw/arm/shoulder pain (especially with shortness of breath, unusual sweating), unexplained weight loss. In addition, before you self-treat with this medication, get medical help right away if you have any of these signs of a serious condition: trouble/pain swallowing food, bloody vomit, vomit that looks like coffee grounds, bloody/black stools, heartburn for over 3 months, frequent chest pain, frequent wheezing (especially with heartburn), nausea/vomiting, stomach pain. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Proton pump inhibitors (such as esomeprazole) may increase your risk for bone fractures, especially with longer use, higher doses, and in older adults. Talk with your doctor or pharmacist about ways to prevent bone loss/fracture, such as by taking calcium (such as calcium citrate) and vitamin D supplements. Older adults

may be more sensitive to the side effects of this drug, especially bone loss and fractures (see above), and C. difficile infection (see Side Effects section). Children may be more sensitive to the side effects of this drug, especially fever, cough, and infections of the nose/throat/airways. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. However, similar drugs pass into breast milk. The effects on a nursing infant are unknown. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: confusion, extreme sweating, blurred vision, or unusually fast heartbeat. If your doctor has prescribed this medication for you, do not share it with others. If your doctor instructs you to use this medication regularly for a long time, lab and/or medical tests (such as a magnesium blood test, vitamin B-12 levels) may be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: clopidogrel, mavacamten, methotrexate (especially high-dose treatment), rifampin, St John's wort. Some products need stomach acid so that the body can absorb them properly. Esomeprazole decreases stomach acid, so it may change how well these products work. Some affected products include atazanavir, erlotinib, levoketoconazole, nelfinavir, pazopanib, rilpivirine, sparsentan, certain azole antifungals (itraconazole, ketoconazole, posaconazole), among others. Esomeprazole is very similar to omeprazole. Do not use any medications containing omeprazole while using esomeprazole. This medication may interfere with certain lab tests, possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug.

Side Effects			
Uses			

Interactions			
Other Details			