## iMedix: Your Personal Health Advisor.

## Niaspan

Search for medical details concerning Niaspan oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- ActiveIngredient:
- DosageForm:
- Dosage:
- Indications:
- Manufacturer:
- Storage:
- Market Price:
- Drug Status:

## **Description**

Side Effects Flushing of the face and neck along with warmth, headache, itching, burning, sweating, chills, or tingling may occur within 2-4 hours of taking this medication. Flushing may last for a few hours after use. These effects should improve or go away as your body adjusts to the medication. Dizziness, stomach upset, heartburn, nausea, vomiting, and diarrhea may also occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. To reduce the risk of dizziness and lightheadedness, get up slowly when rising from a sitting or lying position. This is very important if you are also taking medications to lower your blood pressure. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: nausea/vomiting that doesn't stop, fast/irregular heartbeat, severe headache (migraine), unusual joint pain, swelling of legs/arms, vision problems, severe stomach/abdominal pain, black stools, easy bruising/bleeding, unexplained muscle pain/tenderness/weakness, signs of kidney problems (such as change in the amount of urine), dark urine, vomit that looks like coffee grounds, yellowing eyes/skin. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Niacin is used with a proper diet and exercise program to help lower "bad" cholesterol and fats (LDL, triglycerides) and raise "good" cholesterol (HDL) in the blood. It is generally used after non-drug treatments have not been fully successful at lowering cholesterol. Niacin is also known as vitamin B-3 (nicotinic acid), one of the Bcomplex vitamins. It may be used with or without other medications. Lowering "bad" cholesterol/triglycerides and raising "good" cholesterol helps prevent strokes and heart attacks. Lowering fats may also help reduce the risk of pancreas problems (pancreatitis) in people at risk. In addition to eating a proper diet (such as a low-cholesterol/low-fat diet), other lifestyle changes that may help this medication work better include exercising, losing weight if overweight, and stopping smoking. Consult your doctor for more details. How to use Niaspan Read the Patient Information Leaflet if available from your pharmacist before you start taking niacin and each time you get a refill. If you have any questions, ask your doctor or

pharmacist. Take this medication by mouth with a low-fat meal or snack as directed by your doctor, usually at bedtime. Taking niacin on an empty stomach increases side effects (such as flushing, upset stomach). Swallow the tablets whole. Do not crush or chew the tablets. Doing so can release all of the drug at once, increasing the risk of side effects. Niacin is available in different formulations (such as immediate and sustained release). Do not switch between strengths, brands, or forms of niacin. Severe liver problems may occur. The dosage is based on your medical condition and response to treatment. To reduce your risk of side effects, your doctor may direct you to start this medication at a low dose and gradually increase your dose. Your dose will need to be increased slowly, even if you are already taking niacin and are being switched from another niacin product (such as immediate-release tablets) to this product. Follow your doctor's instructions carefully. If you also take certain other drugs to lower your cholesterol (bile acid-binding resins such as cholestyramine or colestipol), take niacin at least 4-6 hours before or after taking these medications. These products can react with niacin, preventing its full absorption. Continue to take other medications to lower your cholesterol as directed by your doctor. To lessen the chance of side effects such as flushing, avoid alcohol, hot beverages, and eating spicy foods near the time you take niacin. Taking a plain (non-enteric coated, 325 milligram) aspirin or a nonsteroidal anti-inflammatory drug (such as ibuprofen, 200 milligrams) 30 minutes before taking niacin may help prevent flushing. Ask your doctor if this treatment is right for you. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day. Keep taking this medication even if you feel well. Most people with high cholesterol do not feel sick. Do not stop taking this medication without consulting your doctor. If you stop taking niacin, you may need to return to your original dose and gradually increase it again. Ask your doctor or pharmacist for instructions on restarting your dose if you have not taken your medication for an extended period (more than 7 days). It is very important to continue to follow your doctor's advice about diet and exercise. Precautions Before taking niacin, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: very low blood pressure, alcohol use, bleeding problems (such as low platelets), diabetes, gallbladder disease, glaucoma, gout, heart disease (such as recent heart attack, unstable angina), kidney disease, liver disease/increase in liver enzymes, untreated mineral imbalance (low phosphate levels), stomach/intestinal ulcers, underactive thyroid (hypothyroidism). This drug may make you dizzy. Alcohol or marijuana (cannabis) can make you more dizzy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). This medication may rarely make your blood sugar rise, which can cause or worsen diabetes. If you already have diabetes, check your blood sugar regularly as directed and share the results with your doctor. Tell your doctor right away if you have symptoms of high blood sugar such as increased thirst/urination. Your doctor may need to adjust your diabetes medication, exercise program, or diet. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. This medication passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Lab tests (such as blood lipids, blood sugar, liver function) should be done while you are taking this medication. Keep all medical and lab appointments. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Check all prescription and nonprescription medicine labels carefully since vitamins/dietary

supplements may also contain niacin or nicotinamide. These may increase your risk of side effects if taken together. Ask your doctor or pharmacist for more details. This medication may interfere with certain lab tests (including urine or blood catecholamines, copper-based urine glucose tests), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug.
Side Effects
Uses
Interactions
Other Details