

# iMedix: Your Personal Health Advisor.

## Nourianz 40 Mg Tablet Antiparkinson Adjuvant – Adenosine Receptor Antagonist

---

Search for medical details concerning Nourianz oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

Side Effects Dizziness, decreased appetite, or constipation may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: worsening muscle twitching/uncontrollable movements, mental/mood changes (such as agitation, hallucinations), unusual strong urges (such as increased gambling, increased sexual urges). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Istradefylline is used by people with Parkinson's disease taking carbidopa/levodopa to reduce the amount of "off" time (periods of slow movement or stiffness). How to use Nourianz 40 Mg Tablet Antiparkinson Adjuvant – Adenosine Receptor Antagonist Take this medication by mouth with or without food as directed by your doctor, usually once daily. The dosage is based on your medical condition, response to treatment, and other medications you may be taking. Be sure to tell your doctor and pharmacist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Do not increase your dose or take it more often than prescribed. Your condition will not improve any faster, and your risk of side effects will increase. Tell your doctor if this medication stops working well or if your condition gets worse. Precautions Before taking istradefylline, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver disease, mental/mood disorders (such as schizophrenia, bipolar disorder). This drug may make you dizzy. Alcohol or marijuana (cannabis) can make you more dizzy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs,

and herbal products). During pregnancy, this medication should be used only when clearly needed. It may harm an unborn baby. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: mental/mood changes (such as agitation, hallucinations). Do not share this medication with others. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. A product that may interact with this drug is: elacestrant. Other medications can affect the removal of istradefylline from your body, which may affect how istradefylline works. Examples include rifamycins (such as rifabutin, rifampin), St. John's wort, drugs used to treat seizures (such as carbamazepine, phenytoin), among others. Cigarette smoking decreases blood levels of this medication. Tell your doctor if you smoke or if you have recently stopped smoking.

---

---

---

## **Side Effects**

---

---

---

## **Uses**

---

---

---

## **Interactions**

---

---

---

## **Other Details**

