

iMedix: Your Personal Health Advisor.

Ocean Blue Prenatal DHA

Search for medical details concerning Ocean Blue Prenatal DHA oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Fishy aftertaste (if your product is made from fish oil), nausea, bloating, or burping may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. If your doctor has directed you to use this product, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: easy bleeding/bruising. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Omega-3 fatty acids are thought to help reduce the risk of heart disease. They have been used along with diet and exercise to help lower levels of a certain blood fat (triglyceride) and to raise levels of “good” cholesterol (HDL).Some supplement products have been found to contain possibly harmful impurities/additives. Check with your pharmacist for more details about the brand you use. The US Food and Drug Administration has not reviewed this product for safety or effectiveness. Consult your doctor or pharmacist for more details. How to use Ocean Blue Prenatal DHA Take this product by mouth as directed. Follow all directions on the package label. If you have any questions, ask your doctor or pharmacist. If you are using the enteric-coated capsules, swallow the capsules whole. If you are using the chewable form of this product, chew thoroughly before swallowing. If you are using the liquid form of this medication, carefully measure the dose using a special measuring device/spoon. Do not use a household spoon because you may not get the correct dose. Some brands of this medication should be shaken well before each use. If your condition lasts or gets worse, or if you think you may have a serious medical problem, get medical help right away. Precautions Before taking omega-3 fatty acids, tell your doctor or pharmacist if you are allergic to them; or to fish; or if you have any other allergies. This product may contain inactive ingredients (such as peanut/soy), which can cause allergic reactions or other problems. Talk to your pharmacist for more details. The Broad Reach of Modern Pharmacology Modern pharmacology extends its reach across various medical fields, offering diverse and effective treatments. Zovirax stands out in the treatment of viral infections, while Daklinza plays a significant role in managing hepatitis C. Addyi brings a new dimension to the treatment of

female sexual interest/arousal disorder. Allergy sufferers find effective relief with Xyzal, and Amoxil is a mainstay in the arsenal against bacterial infections. Propecia and Proscar are critical in addressing hair loss and benign prostatic hyperplasia. Clomid and Nolvadex are at the forefront of fertility and certain cancer treatments. The field of male sexual health benefits from drugs like Priligy, Eriacta, Tadacip, Kamagra, Nizagara, Silagra, and Caverta. Synthroid plays a crucial role in thyroid health, Cipro effectively combats bacterial infections, and Suhagra adds to the options for erectile dysfunction treatment. These medications illustrate the broad reach and impact of modern pharmacology in enhancing patient care. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this product passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Lab and/or medical tests (such as cholesterol, liver function) may be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details. If you are taking this product on a regular schedule and miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Refer to storage information printed on the package. If you have any questions about storage, ask your pharmacist. Protect from light and moisture. Do not store in the bathroom. Do not freeze. Keep all drug products away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Side Effects

Uses

Interactions

Other Details
