

# iMedix: Your Personal Health Advisor.

## Octreotide Acetate 20 Mg Capsule,Delayed Release

---

Search for medical details concerning octreotide acetate oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

Side Effects Nausea, vomiting, diarrhea, stomach upset, gas, bloating, constipation, dizziness, muscle/joint pain, or headache may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: signs of gallbladder/liver problems (such as fever, stomach/abdominal pain, severe nausea/vomiting, yellowing eyes/skin, unexplained pain in the back/right shoulder), signs of underactive thyroid (such as unexplained weight gain, cold intolerance, slow heartbeat, severe constipation, unusual/extreme tiredness, growth/lump/swelling on the front of the neck), worsening heart condition symptoms (such as trouble breathing, slow/fast/irregular heartbeat), swelling of the arms/legs. This medication may cause changes in blood sugar, especially if you have diabetes. Symptoms of high blood sugar include increased thirst and urination. Symptoms of low blood sugar include sudden sweating, shaking, fast heartbeat, hunger, blurred vision, dizziness, or tingling hands/feet. Tell your doctor right away if you experience symptoms of high or low blood sugar while using this medication. If you already have diabetes, check your blood sugar regularly as directed and share the results with your doctor. Your doctor may need to adjust your diabetes medication, exercise program, or diet. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Octreotide is used to treat a certain condition (acromegaly) that occurs when the body makes too much of a certain natural substance called growth hormone. Treating acromegaly helps reduce the risk of serious problems such as diabetes and heart disease. Octreotide works by decreasing how much growth hormone your body makes. This drug is not a cure for acromegaly. This medication is usually used with other treatment (such as surgery, radiation, other drugs). How to use Octreotide Acetate 20 Mg Capsule,Delayed Release Read the Patient Information Leaflet and Instructions for Use if available from your pharmacist before you start using octreotide and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth on an

empty stomach at least 1 hour before a meal or at least 2 hours after a meal as directed by your doctor, usually twice daily. Swallow the capsule(s) whole with a glass of water. Do not crush or chew the capsules. Doing so can release all of the drug at once, increasing the risk of side effects. The dosage is based on your medical condition, response to treatment, lab tests, and other medications you may be taking. Be sure to tell your doctor and pharmacist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). To reduce your risk of side effects, your doctor may direct you to start this medication at a low dose and gradually increase your dose. Your doctor may also stop your medication from time to time to monitor your medical condition. Follow your doctor's instructions carefully. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same times each day. Tell your doctor if your condition does not get better or if it gets worse. **Precautions** Before using octreotide, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney problems, liver disease (such as cirrhosis), diabetes, thyroid problems, gallbladder problems (such as gallstones), heart problems (such as heart failure), nutrition problems (such as decreased fat absorption, vitamin B12 deficiency). This drug may make you dizzy. Alcohol or marijuana (cannabis) can make you more dizzy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). This medication may restore the normal ability to become pregnant in females with acromegaly who have infertility. Females of childbearing age should discuss reliable forms of birth control with the doctor. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. **Overdose** If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Lab and/or medical tests (such as blood sugar, thyroid function, hormone levels, vitamin B12 levels) should be done before you start using this medication and while you are using it. Keep all medical and lab appointments. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store the unopened medication package (wallet) in the refrigerator. Do not freeze. After opening a medication package, store at room temperature for up to 1 month. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. **Interactions** Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

---

---

## **Side Effects**

---

---

## **Uses**

---

---

---

---

---

## Interactions

---

---

---

---

## Other Details

---

---