

iMedix: Your Personal Health Advisor.

Oritavancin Solution, Reconstituted (Recon Soln)

Search for medical details concerning oritavancin intravenous on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Nausea, vomiting, or headache may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Injection reaction symptoms such as flushing of the upper body, hives, itching, back/chest pain, chills, or shaking may also occur. Tell your doctor right away if you have any of these symptoms. Stopping or slowing down the injection may help. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: bone pain, fast heartbeat, unusual tiredness, swelling ankles/feet/hands. This medication may rarely cause a severe intestinal condition due to a bacteria called *C. difficile*. This condition may occur during treatment or weeks to months after treatment has stopped. Tell your doctor right away if you develop: diarrhea that doesn't stop, abdominal or stomach pain/cramping, blood/mucus in your stool. If you have these symptoms, do not use anti-diarrhea or opioid products because they may make symptoms worse. Use of this medication for prolonged or repeated periods may result in oral thrush or a new yeast infection. Contact your doctor if you notice white patches in your mouth, a change in vaginal discharge, or other new symptoms. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is an antibiotic used to treat serious bacterial infections of the skin. It works by stopping the growth of bacteria. How to use Oritavancin Solution, Reconstituted (Recon Soln) This medication is given by a health care professional as a single dose. Depending on your product, this medication is injected slowly into a vein over 1 or 3 hours. Tell your doctor if your condition lasts or gets worse. Precautions Before using oritavancin, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history. Oritavancin may cause live bacterial vaccines (such as typhoid vaccine) to not work well. Tell your health care professional that you are using oritavancin before having any immunizations/vaccinations. Before having surgery, tell your doctor or dentist

about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Lab and/or medical tests (such as liver function) may be done while you are using this medication. Keep all medical and lab appointments. Consult your doctor for more details. Not applicable. Not applicable. This medication is given in a hospital and will not be stored at home. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. One product that may interact with this drug is: heparin. This medication may interfere with certain lab tests (including blood clotting tests such as PT/INR/aPTT), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug, and if you are using warfarin or are being treated for bleeding/clotting disorders.

Side Effects

Uses

Interactions

Other Details
