

iMedix: Your Personal Health Advisor.

Pedialyte ELECTROLYTE Singles

Search for medical details concerning Pedialyte Singles oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Mild nausea and vomiting may occur. These effects can be decreased by taking this product slowly in small amounts with a spoon. If any of these effects last or get worse, contact your doctor or pharmacist promptly. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: dizziness, unusual weakness, swelling of ankles/feet, mental/mood changes (such as irritability, restlessness), seizures. A very serious allergic reaction to this product is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This product is used to replace fluids and minerals (such as sodium, potassium) lost due to diarrhea and vomiting. It helps prevent or treat the dehydration. Having the right amount of fluids and minerals is important for the normal functioning of the body. How to use Pedialyte ELECTROLYTE Singles Take this product by mouth as directed by your doctor, or follow all directions on the product package. If you have any questions, ask your doctor or pharmacist. Some brands of the liquid form should be mixed before using, while others should not be mixed. Ask your doctor or pharmacist if you have questions about your specific brand. If your liquid form is a suspension, shake the bottle well before each dose. If you are using the powdered form, mix the contents of the packet in water as directed on the product package. Stir until completely dissolved. Do not heat this product unless the product package states that it is okay to do so. Dosage is based on your medical condition and response to treatment. Do not drink fruit juices or eat foods with added salt while taking this product unless directed by your doctor. Tell your doctor if your condition gets worse, if it lasts for more than 24 hours, or if you also have severe stomach/abdominal pain, blood in the stool/vomit, a fever, or signs of dehydration (such as dizziness, decreased urination, severe thirst, very dry mouth, seizures). If you think you may have a serious medical problem, get medical help right away. Precautions Before taking this product, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. If you have

any of the following health problems, consult your doctor or pharmacist before using this product: inability to keep fluids in the stomach (for example, due to vomiting that is severe or doesn't stop), difficulty urinating, intestinal blockage. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Tell your doctor if you are pregnant before using this medication. It is unknown if this product passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Keep all medical and lab appointments. As your condition improves, you can start eating soft, bland foods such as rice, bananas, toast, boiled potatoes, or other foods recommended by your doctor. Not applicable. Refer to storage information printed on the package. If you have any questions about storage, ask your pharmacist. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: mineral supplements (such as potassium).

Side Effects

Uses

Interactions

Other Details
