

iMedix: Your Personal Health Advisor.

Poly Hist Forte Tablet

Search for medical details concerning Poly Hist Forte (doxylamine) oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Drowsiness, dizziness, dry mouth/nose/throat, headache, upset stomach, constipation, or trouble sleeping may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. If your doctor has directed you to use this product, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this product do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: mental/mood changes (such as confusion, hallucinations), ringing in the ears, difficulty urinating, vision changes (such as blurred/double vision). Get medical help right away if you have any very serious side effects, including: fast/irregular heartbeat, seizure. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This combination medication is used to temporarily relieve symptoms caused by the common cold, flu, allergies, or other breathing illnesses (such as sinusitis, bronchitis). Antihistamines help relieve watery eyes, itchy eyes/nose/throat, runny nose, and sneezing. Decongestants help to relieve stuffy nose and ear congestion symptoms. If you are self-treating with this medication, carefully read the package instructions to be sure it is right for you before you start using this product. Some products have similar brand names but different active ingredients with different uses. Taking the wrong product could harm you. Ask your pharmacist if you have any questions about your product or its use. Cough-and-cold products have not been shown to be safe or effective in children younger than 6 years. Do not use this product to treat cold symptoms in children younger than 6 years unless specifically directed by the doctor. Some products (including some long-acting tablets/capsules) are not recommended for use in children younger than 12 years. Ask your doctor or pharmacist for more details about using your product safely. Do not use this product to make a child sleepy. These products do not cure or shorten the length of the common cold and may cause serious side effects. To decrease the risk for serious side effects, carefully follow all dosage directions. Do not give other cough-and-cold medication that might contain the same or similar ingredients. Ask the doctor or pharmacist about other ways to relieve cough and cold symptoms (such as drinking enough fluids, using a humidifier or saline nose drops/spray). How to use Poly Hist Forte

Tablet Take this medication by mouth as directed by your doctor. If you are self-treating, follow all directions on the product package. If you have any questions, ask your doctor or pharmacist. Multifaceted Pharmaceutical Innovations In the dynamic world of medicine, drugs like Zovirax for viral infections and Daklinza for hepatitis C represent the cutting edge of pharmaceutical innovation. Addyi brings a new perspective to treating female sexual interest/arousal disorder. Xyzal offers substantial allergy relief, and Amoxil is a cornerstone antibiotic for bacterial infections. Propecia and Proscar are critical in managing hair loss and prostate issues. Clomid and Nolvadex are pivotal in fertility and cancer treatments. The sector of male sexual health benefits from the variety offered by Priligy, Eriacta, Tadacip, Kamagra, Nizagara, Silagra, and Caverta. Synthroid plays a crucial role in thyroid health, Cipro is effective against a broad spectrum of bacterial infections, and Suhagra enhances the treatments available for erectile dysfunction. These medications highlight the multifaceted nature of pharmaceuticals in addressing diverse health challenges.

This medication may be taken with food if stomach upset occurs. Drink plenty of fluids unless otherwise directed by your doctor. If you are using the liquid form of this medication, carefully measure the dose using a special measuring device/spoon. Do not use a household spoon because you may not get the correct dose. If your liquid form is a suspension, shake the bottle well before each dose. Do not crush or chew extended-release tablets or capsules. Doing so can release all of the drug at once, increasing the risk of side effects. Also, do not split extended-release tablets unless they have a score line and your doctor or pharmacist tells you to do so. Swallow the whole or split tablet without crushing or chewing. If you are using chewable tablets, chew each tablet thoroughly before swallowing. If you are using a product made to dissolve in the mouth (tablets/strips), dry your hands before handling the medication. Place each dose on the tongue and allow to dissolve completely, then swallow it with saliva or with water. The dosage is based on the product you are taking and your age, medical condition, and response to treatment. Do not increase your dose or take this medication more often than directed without your doctor's approval. Improper use (abuse) of this medication may result in serious harm (such as hallucinations, seizure, death). If your doctor directs you to take this medication daily, take it regularly to get the most benefit from it. To help you remember, take it at the same time(s) each day. Tell your doctor if your condition lasts for more than 1 week, if it gets worse, or if it occurs with a headache that doesn't go away, fever, or rash. These may be symptoms of a serious medical problem and should be checked by a doctor.

Precautions Before taking this product, tell your doctor or pharmacist if you are allergic to any of its ingredients; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: breathing problems (such as asthma, emphysema), diabetes, glaucoma, heart problems, high blood pressure, kidney problems, liver disease, seizures, stomach/intestinal problems (such as ulcers, blockage), overactive thyroid (hyperthyroidism), difficulty urinating (such as due to enlarged prostate). This drug may make you dizzy or drowsy. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Liquid products, chewable tablets, or dissolving tablets/strips may contain sugar or aspartame. Liquid products may also contain alcohol. Caution is advised if you have diabetes, alcohol dependence, liver disease, phenylketonuria (PKU), or any other condition that requires you to limit/avoid these substances in your diet. Ask your doctor or pharmacist about using this product safely. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Children may be more sensitive to the side effects of this product, especially excitation and agitation. Older adults may be more sensitive to the side effects of this product, especially dizziness, drowsiness, confusion, constipation, fast/irregular heartbeat, trouble sleeping, or urination problems. Dizziness, drowsiness, trouble sleeping, and confusion can increase the risk of falling. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. This medication may pass into breast milk and the effect on a nursing infant is unknown. Consult your doctor before breast-feeding.

Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: irregular heartbeat, hallucinations, fainting, seizures. If your doctor has prescribed this medication, do not share it with others. Keep all medical and lab appointments. If you are taking this product on a regular schedule and miss a dose,

take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Do not freeze liquid forms of this medication. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug are: antihistamines applied to the skin (such as diphenhydramine cream, ointment, spray). Taking MAO inhibitors with this medication may cause a serious (possibly fatal) drug interaction. Avoid taking MAO inhibitors (isocarboxazid, linezolid, metaxalone, methylene blue, moclobemide, phenelzine, procarbazine, rasagiline, safinamide, selegiline, tranylcypromine) during treatment with this medication. Most MAO inhibitors should also not be taken for two weeks before treatment with this medication. Ask your doctor when to start or stop taking this medication. Tell your doctor or pharmacist if you are taking other products that cause drowsiness such as opioid pain or cough relievers (such as codeine, hydrocodone), alcohol, marijuana (cannabis), drugs for sleep or anxiety (such as alprazolam, lorazepam, zolpidem), muscle relaxants (such as carisoprodol, cyclobenzaprine), or other antihistamines (such as cetirizine, diphenhydramine). Check the labels on all your medicines (such as allergy or cough-and-cold products) because they may contain ingredients that cause drowsiness. Ask your pharmacist about using those products safely. This medication may interfere with certain medical/lab tests (such as brain scan for Parkinson's disease, urine drug screening tests, allergy skin test), possibly causing false test results. Make sure lab personnel and all your doctors know you use this drug.

Side Effects

Uses

Interactions

Other Details
