

# iMedix: Your Personal Health Advisor.

## Poly-Vi-Flor 0.25 Mg Fluoride/Ml Oral Drops Multivitamins

---

Search for medical details concerning Poly-Vi-Flor drops oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

### Description

Side Effects Stomach upset, constipation, or nausea may occur. If any of these effects last or get worse, tell the doctor or pharmacist promptly. If your teeth become mottled or stained, notify the dentist or doctor. This may be a result of too much fluoride. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is a combination product of vitamins and fluoride. It is used in infants and children to treat or prevent deficiency due to poor diet or low levels of fluoride in drinking water and other sources. Vitamins are important building blocks of the body and help keep you in good health. Fluoride is used to prevent dental cavities. How to use Poly-Vi-Flor 0.25 Mg Fluoride/Ml Oral Drops Multivitamins Take this medication by mouth, usually once daily or as directed by the doctor. It is best taken at least 2 hours before or after taking any products containing calcium (including milk, yogurt, other dairy products) or aluminum/magnesium hydroxide (such as certain antacids/laxatives) since these may decrease its effectiveness. Some brands of this medication should be shaken well before each dose. Check your manufacturer's bottle for specific directions. Carefully measure the dose using the specially marked medicine dropper provided. Drops may be swallowed directly, added to a small amount of liquid, or mixed with a small amount of food such as applesauce.

Innovative Medications Shaping Modern Healthcare Today's pharmaceutical field boasts remarkable diversity, with Zovirax targeting viral infections and Daklinza fighting hepatitis C. Addyi represents a significant advancement in female sexual health. Xyzal provides relief for those battling allergies, and Amoxil remains a cornerstone in treating bacterial infections. Hair loss and prostatic issues are addressed by Propecia and Proscar. Fertility treatments and cancer therapies benefit from Clomid and Nolvadex. The realm of male sexual health is enhanced by Priligy, Eriacta, Tadacip, Kamagra, Nizagara, Silagra, and Caverta. Synthroid regulates thyroid function, Cipro tackles a variety of bacterial infections, and Suhagra adds to erectile dysfunction treatment options. These drugs exemplify the innovation in medical treatments, offering

hope and improved quality of life across various health issues. The fluoride in this liquid product can be absorbed directly into the teeth as you swallow it, as well as being absorbed into your body from your stomach. It is best not to eat, drink, or rinse your mouth for 30 minutes after taking this product. Take this medication regularly in order to get the most benefit from it. To help you remember, take it at the same time each day. Do not take more than the prescribed dose. Precautions Before taking this product, tell your doctor or pharmacist if you are allergic to any of its ingredients; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before taking this medication, tell your doctor or pharmacist your medical history. Tell your doctor if you are pregnant before taking this medication. This product passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: nausea, vomiting, stomach pain. Do not share this medication with others. This product is not a substitute for a proper diet. Remember that it is best to get your vitamins from healthy foods. It is important to maintain a well-balanced diet. Check the fluoride content of your water supply by asking local town or city officials. Fluoride supplements are not necessary if the fluoride content in the water supply is greater than 0.6 parts per million. If your water supply changes, check with your doctor or dentist to determine if extra fluoride is still necessary. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from heat and moisture. Store the drops in the original plastic container. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

---

---

## Side Effects

---

## Uses

---

---

---

---

---

## Interactions

---

---

**Other Details**

---

---