

iMedix: Your Personal Health Advisor.

Generic Priligy (Dapoxetine)

Priligy (Dapoxetine) is a novel medication specifically designed for the treatment of premature ejaculation in adult men. Priligy is designed for short-term use before sexual activity. Priligy offers a significant improvement in the control over ejaculation.

- **ActiveIngredient:** Dapoxetine Hydrochloride
 - **DosageForm:** Tablets
 - **Dosage:** 30 mg
 - **Indications:** Premature Ejaculation (PE)
 - **Manufacturer:** Sunrise Remedies Pvt. Ltd.
 - **Storage:** Store in a cool, dry place away from direct sunlight and moisture. Keep out of reach of children.
 - **Market Price:** 0.95
 - **Drug Status:** Prescription Only
-

Description

What is Priligy? Priligy (Dapoxetine) is a medication used to treat premature ejaculation (PE), a common sexual health condition in men. It contains the active ingredient dapoxetine, which is a short-acting selective serotonin reuptake inhibitor (SSRI). Dapoxetine helps delay ejaculation by increasing the levels of serotonin in the brain, which aids in controlling the ejaculation reflex. How Priligy Works Priligy works by boosting serotonin levels in the brain, enhancing the ability to delay ejaculation. This increase in serotonin helps control the timing of ejaculation, providing better control for those experiencing PE. Priligy's Availability Priligy (Dapoxetine) is a prescription medication, meaning it should only be obtained with a prescription from a qualified healthcare professional. It's crucial to get this medication from reputable pharmacies to ensure its safety and efficacy.

General Instructions

How to Take Priligy Generic Priligy is typically taken as one tablet 1-3 hours before sexual activity. It's important to swallow the tablet with water. The usual starting dose is 30mg, which may be increased to 60mg based on individual response and tolerance. Remember, Dapoxetine (Priligy) is not meant for continuous daily use. Important Precautions Medication Interactions: Priligy can interact with other medications. Always inform your doctor about any other drugs you are taking, including over-the-counter medications and supplements. Consult Your Doctor: Before starting Priligy, consult your healthcare provider to ensure it's safe for you, especially if you have any underlying health conditions. Do Not Exceed Recommended Dosages: Taking more than the recommended dose can increase the risk of side effects and should be avoided.

Side Effects

Common Side Effects Priligy (Dapoxetine) may cause some common side effects, which include: Nausea: Feeling sick to your stomach or like you might vomit. Dizziness: Feeling lightheaded, unsteady, or like you might faint. Headache: Pain or discomfort in the head or face area. Diarrhea: Loose or watery bowel movements. Runny Nose: Excess mucus discharge from the nose, often caused by irritation or infection. These side effects are usually mild and go away on their own as your body gets used to the medication.

Serious Side Effects While rare, some serious side effects may occur: Seizures: Sudden, uncontrolled electrical activity in the brain, which can cause changes in behavior, movements, or consciousness. Serotonin Syndrome: A potentially life-threatening condition that occurs when there's too much serotonin in the body. Symptoms can include agitation, confusion, rapid heart rate, high blood pressure, dilated pupils, and muscle rigidity. Allergic Reactions: Severe allergic reactions can include symptoms such as rash, itching, swelling, severe dizziness, and trouble breathing. If you experience any of these serious side effects, seek immediate medical attention.

Uses

Treating Premature Ejaculation Priligy is used to help men who suffer from premature ejaculation (PE), which is when ejaculation happens sooner during sexual activity than a man or his partner would like. Generic Priligy helps delay ejaculation and improve control over ejaculation, which can lead to longer-lasting and more satisfying sexual experiences. It's important to note that Dapoxetine (Priligy) is not a cure for PE but a treatment to manage the condition and improve sexual function and satisfaction.

Safety advice

Interactions Alcohol:

- Unsafe
- Combining Priligy 30 mg with alcohol can increase the risk of side effects like dizziness, fainting, and impaired judgment. It's recommended to avoid alcohol when taking this medication.

Interactions Other Medications:

- Consult your doctor
- Priligy 30 mg can interact with other medications, particularly antidepressants, certain painkillers, and medications for heart conditions. Always consult your doctor before combining Priligy with other drugs.

Special Groups Pregnancy:

- Unsafe
- Priligy 30 mg is not intended for use by women, including those who are pregnant. It is specifically formulated for men with premature ejaculation.

Special Groups Breastfeeding:

- Unsafe
- This medication should not be used by breastfeeding women as it is designed exclusively for men and may pose risks to infants.

Special Groups Elderly:

- Consult your doctor
- Priligy 30 mg is generally not recommended for elderly patients due to limited data on its safety and efficacy in this age group. A doctor's consultation is advised.

Special Groups Children:

- Unsafe

- Priligy 30 mg is not suitable for individuals under 18 years of age and should not be used by children or adolescents.

Effects on Activities Driving:

- Use with caution
- Priligy 30 mg may cause dizziness, fainting, or blurred vision, which can impair your ability to drive safely. Use caution and avoid driving if you feel unwell.

Effects on Activities Operating Machinery:

- Use with caution
 - Operating machinery should be done with caution when taking Priligy 30 mg, as the medication can affect alertness and coordination.
-

Concerns

Safety and Effectiveness Priligy has been shown to be effective for many men with premature ejaculation, but it is not a guaranteed solution for everyone. Its effectiveness can vary based on individual differences and the underlying causes of PE. Some men may find significant improvement, while others may see less benefit.

Potential for Dependence While Priligy (Dapoxetine) is not considered addictive, there is a potential concern about relying on it for long-term use. It's important to discuss any long-term use of Priligy with your doctor to ensure it's safe and appropriate for your situation.

When Not to Use Conditions that May Prevent Use Priligy should not be used by individuals with certain medical conditions, such as: Severe Liver or Kidney Disease: These conditions can affect how the body processes the medication. Certain Heart Conditions: Such as significant heart disease or a history of fainting. Allergies to Dapoxetine: If you have a known allergy to the active ingredient in Priligy or any other component of the formulation. If you have any of these conditions, it's important to consult your doctor before using Dapoxetine (Priligy). They can help determine if this medication is safe for you to use.

Warnings

Alcohol Avoid drinking alcohol while taking Priligy, as it can make side effects like dizziness and fainting worse. This combination increases the risk of accidents or injuries because you might feel very lightheaded or unsteady.

Other Medications Priligy can interact with certain medications, making side effects more likely or reducing the effectiveness of Priligy. These include: Monoamine Oxidase Inhibitors (MAOIs): Medications used to treat depression and other mental health conditions. Certain Antidepressants: Such as selective serotonin reuptake inhibitors (SSRIs) and serotonin-norepinephrine reuptake inhibitors (SNRIs), which are also used for treating depression. Some Antifungal Medications: Drugs used to treat fungal infections. Always tell your doctor about all the medications you are taking, including those bought without a prescription and supplements, before starting Generic Priligy. This helps avoid harmful interactions.

Driving and Operating Machinery Priligy (Dapoxetine) can make you feel dizzy or affect your concentration. Until you know how Dapoxetine (Priligy) affects you, avoid driving or using heavy machinery. This is to prevent accidents caused by feeling lightheaded or having difficulty focusing.

Dosage

Typical Dosage The recommended starting dose of Priligy is 30mg, taken as needed approximately 1 to 3 hours before sexual activity. If the 30mg dose is not effective and the side effects are manageable, the dose may be increased to 60mg. The maximum recommended dosing frequency is once every 24 hours.

Important Notes Follow Doctor's Instructions: Always follow your doctor's instructions regarding the dosage. Do not adjust the dose on your own. Do Not Exceed Recommended Dose: Taking more than the recommended dose

can increase the risk of side effects.

Interactions

Medications Priligy (Dapoxetine) can interact with certain medications, leading to increased risk of side effects or reduced effectiveness. These medications include: Monoamine Oxidase Inhibitors (MAOIs): Drugs used to treat depression and other mental health conditions. Taking Generic Priligy with MAOIs can increase the risk of serious side effects. Certain Antidepressants: Including selective serotonin reuptake inhibitors (SSRIs) and serotonin-norepinephrine reuptake inhibitors (SNRIs). Combining these with Priligy can lead to an excessive increase in serotonin levels, which can be dangerous. Some Antifungal Medications: These drugs can affect how Dapoxetine (Priligy) is metabolized in the body, potentially leading to higher levels of the medication and increased side effects. Always inform your healthcare provider about all the medications you are taking, including prescription drugs, over-the-counter medications, and supplements, to avoid harmful interactions.

FAQs

- **Can I take Priligy 30 mg every day?**
Priligy 30 mg is not intended for daily use. It should only be taken when you plan to have sex. Always follow your doctor's instructions on how often to use this medication.
 - **Does Priligy 30 mg affect fertility?**
Priligy 30 mg does not affect fertility. It works by altering the chemicals in the brain to help delay ejaculation and does not impact sperm quality or your ability to father a child.
 - **Can I take Priligy 30 mg if I have heart problems?**
Priligy 30 mg may not be suitable for people with certain heart conditions. You should always discuss your medical history with your doctor to ensure this medication is safe for you.
 - **How long does Priligy 30 mg last?**
The effects of Priligy 30 mg usually last for about 4-6 hours. This can vary from person to person, depending on factors like age, metabolism, and overall health.
-

Other Details

Food Generic Priligy can be taken with or without food. Taking it with food might help reduce the likelihood of stomach upset, which can be a side effect for some people. Duration of Effect The effects of Priligy typically last for several hours. Most men find that Priligy helps them delay ejaculation for the duration of a sexual activity session. Who can use Priligy? Priligy is prescribed for adult men aged 18 to 64 who have all of the following: a history of premature ejaculation, poor control over ejaculation, and a marked personal distress or interpersonal difficulty as a result of PE. Is a prescription required for Priligy? Yes, Priligy is a prescription medication. It's important to consult with a healthcare provider to ensure it's safe and suitable for you to use. Can Priligy cure premature ejaculation permanently? Priligy is not a cure for premature ejaculation but a treatment to manage the condition. It helps to prolong the time taken to ejaculate and can improve control over ejaculation. Can Priligy be taken with other medications? Priligy can interact with certain medications, so it's important to inform your healthcare provider about all medications you are currently taking, including over-the-counter drugs and supplements. Conclusion Seeking Medical Advice It's essential to consult with a healthcare professional before starting Dapoxetine (Priligy) to ensure it is safe and appropriate for you. A doctor can provide personalized advice based on your medical history and current medications. Responsible Use of Medication Use Priligy only as prescribed by your healthcare provider. Do not share it with others or use it for conditions it was not prescribed for. Always follow the dosage

instructions provided by your doctor.

References

Patient.info: Health Information and Symptom Checker. Dapoxetine tablets. In: Patient.info. 2022. [Accessed 23 May 2024]. Available from: <https://patient.info/medicine/dapoxetine-tablets-priligy> National Center for Biotechnology Information. Dapoxetine (Priligy): on demand treatment of premature ejaculation. In: PubMed 2013. [Accessed 23 May 2024]. Available from: <https://pubmed.ncbi.nlm.nih.gov/24564032/> WebMD LLC. Dapoxetine Tablet – Uses, Side Effects, and More: WebMD. 2024. [Accessed 23 May 2024]. Available from: <https://www.webmd.com/drugs/2/drug-91490-2114/duloxetine-oral/duloxetine-sprinkle-capsule-oral/details> Drugs.com. Dapoxetine. In: Drugs.com. 2024. [Accessed 23 May 2024]. Available from: <https://www.drugs.com/international/dapoxetine.html> NPS MedicineWise. Priligy – Dapoxetine. In: nps.org.au; 2024. [Accessed 23 May 2024]. Available from: <https://www.nps.org.au/medicine-finder/priligy-tablets>
