

iMedix: Your Personal Health Advisor.

Rapid Clear Pads, Medicated

Search for medical details concerning Rapid Clear Treatment Pads topical on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Skin reactions such as peeling/burning/dry/reddened skin may occur, especially at the start of treatment. If any of these effects last or get worse, tell your doctor or pharmacist promptly. You may need to apply smaller amounts of the drug or use it less often. Consult your doctor or pharmacist for more details. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This product is used to treat mild to moderate acne. It may be used in combination with other acne treatments. Check with your doctor or pharmacist first before combining acne treatments. Salicylic acid belongs to a class of drugs known as salicylates. When applied to the skin, salicylic acid may work by helping the skin to shed dead cells from the top layer and by decreasing redness and swelling (inflammation). This decreases the number of pimples that form and speeds healing. How to use Rapid Clear Pads, Medicated Follow all directions on the product package. If you have any questions, ask your doctor or pharmacist. Before using this product for the first time, check if you are allergic to it. To check, put a small amount on one or two small areas of acne for three days. If you have an allergic reaction, stop using this product and get medical help right away (see the symptoms of an allergic reaction in the Side Effects section). If you don't have an allergic reaction, you can use the product. This medication is for use on the skin only. To avoid irritation, do not let this medication come into contact with your eyes, nose, mouth, groin, or any broken skin (such as cuts, scrapes, rashes). If you do get the medication in those areas, flush the area with cool water for 15 minutes. If you are using a cream, lotion, solution, or gel, apply a thin film of the medication to the areas of your skin affected by acne, usually 1 to 2 times a day or as directed by your doctor. Before applying each dose, gently wash the affected area with a mild cleanser, then pat dry. Apply a small amount of medicine to the affected area and rub in gently. If using the pads, wash the affected skin with a mild cleanser, then pat dry. Wipe the pad gently onto the skin to apply the medication, usually 1 to 3 times a day or as directed by your doctor. Wash hands after use. If using cleansers containing salicylic

acid, wet the affected area. Gently rub the cleanser into the skin for 10 to 20 seconds. Do not scrub the skin. Work into a full lather and rinse thoroughly, and then pat dry. If too much drying occurs, you may need to rinse the cleanser off sooner or use it less often. Consult your doctor or pharmacist for further details. Dosage is based on your medical condition and response to treatment. Do not use large amounts, apply this drug more often, or use it for a longer period than directed. Your condition will not clear faster, but the chance for side effects may be increased. If your condition lasts or gets worse, or if you think you may have a serious medical problem, get medical help right away. Precautions Before using salicylic acid, tell your doctor or pharmacist if you are allergic to it; or to nonsteroidal anti-inflammatory drugs-NSAIDs (such as aspirin, ibuprofen, naproxen); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. If you have any of the following health problems, consult your doctor or pharmacist before using this product: diabetes, poor circulation, skin problems (such as infection, wounds, atopic dermatitis). During pregnancy, this product should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. Consult your doctor before breast-feeding. Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. If this medication has been prescribed by your doctor, do not share this medication with others. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature. Different brands of this medication have different storage needs. Check the product package for instructions on how to store your brand, or ask your pharmacist. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Side Effects

Uses

Interactions

Other Details
