

# iMedix: Your Personal Health Advisor.

## Ricola 8 Mg Lozenges Not Applicable

---

Search for medical details concerning Ricola mucous membrane on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

**Side Effects** Numbness of the mouth or throat may occur. If this effect lasts or gets worse, tell your doctor or pharmacist promptly. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** This product is used to temporarily help relieve symptoms such as sore throat, throat irritation, or cough (due to a cold, for example). It works by providing a cooling feeling and increasing saliva in the mouth. **Check the ingredients on the label even if you have used the product before.** The manufacturer may have changed the ingredients. Also, products with similar names may contain different ingredients meant for different purposes. **How to use Ricola 8 Mg Lozenges Not Applicable** Read and follow all directions on the product package before using this medication. If you have any questions, consult your pharmacist. If your doctor has directed you to use this product, use it as directed. Let the lozenge dissolve slowly in your mouth and swallow the dissolved liquid along with your saliva. Do not chew or swallow it whole. This product is usually used every 2 hours as needed. If your doctor has directed you to use it, follow your doctor's instructions on how to use it. Do not increase your dose or use this drug more often or for longer than directed. If you have a sore throat that is severe or that lasts more than 2 days, or a cough that lasts more than 7 days or keeps coming back, or symptoms occur with a fever, headache, rash, swelling, nausea, or vomiting, tell your doctor promptly. If sore mouth/throat or cough symptoms do not get better or last more than 7 days, see your doctor or dentist promptly. These could be signs of a serious condition. If you think you may have a serious medical problem, get medical help right away. **Precautions** Before using this product, tell your doctor or pharmacist if you are allergic to menthol; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this product, tell your doctor or pharmacist your medical history, especially of: breathing problems (such as asthma, emphysema), cough with a lot of mucus. This product may contain sugar and/or aspartame. Caution is advised if you have

diabetes, phenylketonuria (PKU), or any other condition that requires you to limit/avoid these substances in your diet. Ask your doctor or pharmacist about using this product safely. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Tell your doctor or pharmacist if you are pregnant before using this product. Menthol passes into breast milk in small amounts. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Keep all regular medical and lab appointments. Not applicable. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

---

---

## **Side Effects**

---

---

## **Uses**

---

---

---

---

---

---

---

## **Interactions**

---

---

## **Other Details**

---

---