

# iMedix: Your Personal Health Advisor.

## Rifadin 600 Mg Intravenous Solution Rifamycins

---

Search for medical details concerning Rifadin intravenous on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

Side Effects Upset stomach, heartburn, nausea, or headache may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. This medication may cause urine, sweat, saliva, or tears to change color (yellow, orange, red, or brown). This effect is harmless and will disappear when the medication is stopped. However, teeth and contact lens staining may be permanent. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: signs of kidney problems (such as change in the amount of urine), mental/mood changes (such as confusion, unusual behavior), unusual tiredness, easy bruising/bleeding, small red spots on the skin, joint pain/swelling, new or worsening shortness of breath. Get medical help right away if you have any very serious side effects, including: chest pain. Rifampin may rarely cause serious (possibly fatal) liver disease. Get medical help right away if you have any symptoms of liver damage, such as: nausea/vomiting that doesn't stop, loss of appetite, stomach/abdominal pain, yellowing eyes/skin, dark urine. This medication may rarely cause a severe intestinal condition due to a bacteria called *C. difficile*. This condition may occur during treatment or weeks to months after treatment has stopped. Tell your doctor right away if you develop: diarrhea that doesn't stop, abdominal or stomach pain/cramping, blood/mucus in your stool. If you have these symptoms, do not use anti-diarrhea or opioid products because they may make symptoms worse. This medication may sometimes cause oral thrush or a new yeast infection. Contact your doctor if you notice white patches in your mouth, a change in vaginal discharge, or other new symptoms. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: fever that doesn't go away, new or worsening lymph node swelling, rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used to prevent and treat tuberculosis and other bacterial infections. Rifampin belongs to a class of drugs known as rifamycin antibiotics. It works by stopping the growth of bacteria. This antibiotic treats only bacterial infections. It will not work for viral infections (such as common cold, flu). Using any antibiotic when it is not needed can cause it to not work for future infections. How to use Rifadin 600 Mg

**Intravenous Solution Rifamycins** This medication is given by injection into a vein as directed by your doctor, usually once or twice daily. It is injected slowly into a vein over 30 minutes. If you are using this medication at home, learn all preparation and usage instructions from your health care professional. Before using, check this product visually for particles or discoloration. If either is present, do not use the liquid. Learn how to store and discard medical supplies safely. The dosage and length of treatment are based on your medical condition, weight, and response to treatment. It is very important to keep using this medication (and other medications used to treat tuberculosis) exactly as prescribed by your doctor. Do not skip any doses. For the best effect, use this antibiotic at evenly spaced times. To help you remember, use this medication at the same time(s) every day. Continue to use this medication until the full prescribed amount is finished, even if symptoms disappear after a few days. Stopping the medication too early may result in a return of the infection. Tell your doctor if your condition lasts or gets worse.

**Precautions** Before using rifampin, tell your doctor or pharmacist if you are allergic to it; or other rifamycins (such as rifabutin); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: diabetes, liver problems (such as hepatitis), HIV infection, history of alcohol use/abuse. Alcohol may increase the risk of liver disease. Avoid alcoholic beverages while using this medication. Rifampin may cause live bacterial vaccines (such as typhoid vaccine) to not work well. Tell your health care professional that you are using rifampin before having any immunizations/vaccinations. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. When this drug is used during the last few weeks of pregnancy, the risk of bleeding in both mother and infant may be increased. Tell your doctor right away if you notice any bleeding in your newborn. Discuss the risks and benefits with your doctor. Rifampin passes into breast milk but is unlikely to harm a nursing infant. Consult your doctor before breast-feeding.

**Overdose** If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Lab and/or medical tests (such as liver/kidney function, bilirubin, complete blood count) may be done before you start using this medication and while you are using it. Keep all medical and lab appointments. Consult your doctor for more details. This medication has been prescribed for your current condition only. Do not use it later for another infection unless your doctor tells you to. It is important to get each dose of this medication as scheduled. If you miss a dose, ask your doctor or pharmacist right away for a new dosing schedule. Do not double the dose to catch up. Consult the product instructions and your pharmacist for storage details. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

**Interactions** Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Rifampin interacts with many medications. While you are using rifampin, it is very important to tell your doctor or pharmacist of any changes in medications that you are taking. Rifampin can speed up the removal of other medications from your body, which may affect how they work. Examples of affected drugs include atovaquone, certain anti-cancer drugs (such as irinotecan), certain azole antifungals (such as itraconazole), certain calcium channel blockers (such as nifedipine), certain products used to treat chronic hepatitis C (such as sofosbuvir), cobicistat, lurasidone, HIV NNRTIs (such as etravirine, nevirapine), praziquantel, HIV protease inhibitors (such as atazanavir, darunavir), among others. This medication may decrease the effectiveness of hormonal birth control such as pills, patch, or ring. This could cause pregnancy. Discuss with your doctor or pharmacist if you should use reliable backup birth control methods while using this medication. Also tell your doctor if you have any new spotting or breakthrough bleeding, because these may be signs that your birth control is not working well. This medication may interfere with certain lab tests (such as urine screening for opiates, blood serum folate/vitamin B12), possibly causing false test results. Rifampin may also make it harder for your body to get rid of certain chemicals used in gallbladder tests. Make sure lab personnel and all your

doctors know you use this drug.

---

---

## Side Effects

---

## Uses

---

---

---

---

---

---

---

## Interactions

---

---

## Other Details

---

---