

iMedix: Your Personal Health Advisor.

Roflumilast 0.3 % Topical Cream

Search for medical details concerning roflumilast topical on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Diarrhea, nausea, headache, or pain at the application site may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: trouble sleeping, signs of infection (such as sore throat that doesn't go away, fever, chills). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, severe itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. How to use Roflumilast 0.3 % Topical Cream Read the Patient Information Leaflet if available from your pharmacist before you start using roflumilast and each time you get a refill. If you have any questions, ask your doctor or pharmacist. This medication is for use on the skin only. Apply this medication to the affected area and rub in completely as directed by your doctor, usually once daily. Wash your hands after applying, unless you are treating your hands. Avoid getting this medication in your eyes, mouth, or vagina. If this happens, rinse right away with plenty of water. Use this medication regularly to get the most benefit from it. To help you remember, use it at the same time each day. Tell your doctor if your condition does not get better or if it gets worse. Precautions Before using roflumilast, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: liver disease. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. To avoid direct contact, do not apply this medication to the nipple and areola if breast-feeding. Consult your doctor before breast-feeding. Consult your pharmacist or physician. Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents

can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. If you miss a dose, apply it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Apply your next dose at the regular time. Do not apply extra medicine to make up for the missed dose. Store at room temperature. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Side Effects

Uses

Interactions

Other Details