

# iMedix: Your Personal Health Advisor.

## Skin Treatment Lotion

---

Search for medical details concerning Skin Treatment topical on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

### Description

Side Effects Stinging, burning, and redness may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: darkening/lightening of the skin, small red dots on the skin. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This medication is used to treat dry, scaly skin conditions (such as xerosis, ichthyosis vulgaris) and can also help relieve itching from these conditions. This medication works by increasing the moisture in the skin. How to use Skin Treatment Lotion If you are using the lotion, shake the bottle well before using. Apply a thin layer of this medication to the affected areas of the skin, usually twice daily or as directed by your doctor. Rub the medication thoroughly into the skin. Be careful to avoid your eyes, lips, inside your mouth/nose, the vaginal area, and any areas of broken skin. Stinging or burning may occur when this medication is applied on the face, on chapped skin, or on a recently shaved area of the skin. Consult your doctor or pharmacist for more details. Inform your doctor if your skin condition does not improve or worsens. Precautions Before using ammonium lactate, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: wounds or sores on the skin. This medication may make you more sensitive to the sun. Limit your time in the sun. Avoid tanning booths and sunlamps. Use sunscreen and wear protective clothing when outdoors. Tell your doctor right away if you get sunburned or have skin blisters/redness. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. Consult your doctor before breast-feeding. Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right

away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Some ways to help prevent dry skin include using lukewarm (not hot) water when bathing, taking baths/showers less often (such as every 1-2 days), keeping baths/showers short, and using a humidifier when the air is very dry. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

---

---

**Side Effects**

---

---

**Uses**

---

---

---

---

---

---

---

**Interactions**

---

---

**Other Details**

---

---