

iMedix: Your Personal Health Advisor.

Sodium Edecrin Vial

Search for medical details concerning Sodium Edecrin intravenous on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Pain at the injection site, dizziness, lightheadedness, weakness, muscle cramps, upset stomach, or diarrhea may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. To reduce the risk of dizziness and lightheadedness, get up slowly when rising from a sitting or lying position. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. This drug can cause dehydration and electrolyte imbalance. Tell your doctor right away if you notice any symptoms of dehydration, such as confusion, unusual decreased urination, unusual dry mouth/thirst, fast/irregular heartbeat, or severe dizziness/lightheadedness. Tell your doctor right away if you have any serious side effects, including: easy bleeding/bruising, feeling of spinning (vertigo), hearing changes (such as ringing or fullness in the ears, decreased hearing/deafness), black/bloody stools, stomach/abdominal pain, vomit that looks like coffee grounds, signs of liver problems (such as nausea/vomiting that doesn't stop, loss of appetite, yellowing eyes/skin, dark urine). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** Ethacrynat sodium is used to decrease swelling (edema) caused by conditions such as cancer, congestive heart failure, liver disease, and kidney disease. This medication works in your kidneys like a “water pill”/diuretic to increase the amount of urine you make. This effect can help your kidneys work better and lessen symptoms such as trouble breathing and swelling in your ankles, feet, hands, or belly. This medication should not be used for infants. **How to use Sodium Edecrin Vial** This medication is given by injection into a vein as directed by your doctor. It is usually given by a health care professional. The dosage is based on your medical condition and response to treatment. Your doctor will adjust your dose based on your medical condition, response to treatment, and lab tests (such as sodium, potassium, chloride levels). If you use this drug too close to bedtime, you may need to wake up to urinate. It is best to use this medication at least 4 hours before your bedtime. Consult your doctor or pharmacist if you have questions about your dosing schedule. Tell your doctor if you do not get better or if you get worse. **Precautions** Before using ethacrynat

sodium, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: gout, kidney disease, liver disease. This drug may make you dizzy. Alcohol or marijuana (cannabis) can make you more dizzy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Avoid alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). This drug can lower certain salt/mineral levels (such as sodium, potassium) in your blood. Your doctor may tell you to use more salt, eat potassium-rich foods (such as bananas, orange juice), or take a potassium supplement. Ask your doctor for more details. This medication may affect your blood sugar. If you have diabetes, check your blood sugar regularly as directed and share the results with your doctor. Your doctor may need to adjust your diabetes medication, exercise program, or diet. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Older adults may be more sensitive to the side effects of this drug, especially dizziness and water/mineral loss. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Lab and/or medical tests (such as kidney tests, blood mineral levels such as potassium) should be done while you are taking this medication. Keep all medical and lab appointments. Not applicable. Not applicable. This medication is given in a hospital or clinic and will not be stored at home. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: desmopressin, furosemide, lithium. Some products have ingredients that could raise your blood pressure or worsen your swelling. Tell your pharmacist what products you are using, and ask how to use them safely (especially cough-and-cold products, diet aids, or NSAIDs such as ibuprofen/naproxen).

Side Effects

Uses

Interactions

Other Details
