

# iMedix: Your Personal Health Advisor.

## Spevigo 60 Mg/Ml Intravenous Solution Antipsoriatics

---

Search for medical details concerning Spevigo intravenous on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

### Description

**Side Effects** Headache, nausea, vomiting, tiredness, or weakness may occur. Pain, redness, swelling, or bruising at the injection site may also occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. This medication may lower your ability to fight infections. This may make you more likely to get a serious (rarely fatal) infection or make any infection you have worse. Get medical help right away if you have any signs of infection (such as sore throat that doesn't go away, fever, swollen lymph nodes, chills, cough). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** This medication is used to treat a severe form of psoriasis (generalized pustular psoriasis). Spesolimab belongs to a class of drugs known as monoclonal antibodies. It works by blocking a certain natural protein in your body (interleukin-36) that may cause inflammation and swelling. **How to use Spevigo 60 Mg/Ml Intravenous Solution Antipsoriatics** Read the Medication Guide provided by your pharmacist before you start using spesolimab and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Before treatment with this medication, your doctor will test you for tuberculosis (TB). Your doctor should also monitor you for symptoms of TB during and after treatment with this drug. If needed, treatment for TB or other infections should be given before using this medication. This medication is given by injection into a vein as directed by your doctor, usually as a single dose injected over 90 minutes. If your symptoms continue, your doctor may prescribe a second dose, usually 1 week after the first dose. The dosage and treatment schedule are based on your medical condition and response to treatment. Tell your doctor if your condition does not get better or if it gets worse.

**Precautions** Before using spesolimab, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: current/recent/returning infection (such as tuberculosis,

herpes). Spesolimab can make you more likely to get infections or may make current infections worse. Stay away from anyone who has an infection that may easily spread (such as chickenpox, COVID-19, measles, flu). Talk to your doctor if you have been exposed to an infection or for more details. Tell your health care professional that you are using spesolimab before having any immunizations/vaccinations. Avoid contact with people who have recently received live vaccines (such as flu vaccine inhaled through the nose). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Keep all medical and lab appointments. Not applicable. Not applicable. This medication is given in a hospital or clinic or doctor's office and will not be stored at home. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

---

---

## **Side Effects**

---

## **Uses**

---

---

---

---

---

---

---

## **Interactions**

---

---

## **Other Details**

---

---

