

# iMedix: Your Personal Health Advisor.

## Super Vidalista

---

Super Vidalista is a combination therapy tablet integrating tadalafil to manage erectile difficulties and dapoxetine to help delay ejaculation. This dual-action formulation aims to address both ED and premature ejaculation concurrently in a single dose for adult men.

- **ActiveIngredient:** Tadalafil, Dapoxetine
  - **DosageForm:** Oral Tablet
  - **Dosage:** 20 mg Tadalafil / 60 mg Dapoxetine per tablet
  - **Indications:** Simultaneous treatment of erectile dysfunction (ED) and premature ejaculation (PE) in adult men
  - **Manufacturer:** Varies (Centurion Laboratories is often cited; verify on packaging)
  - **Storage:** Store below 30°C (86°F) in a dry place, shielded from light and moisture. Keep in original packaging until use. Ensure it is inaccessible to children.
  - **Market Price:**
  - **Drug Status:** Prescription Only
- 

## Description

What is Super Vidalista? Super Vidalista is a composite medication formulated to tackle two distinct male sexual health issues simultaneously. Each tablet contains a precise combination of two active substances: Tadalafil (20 mg): A phosphodiesterase type 5 (PDE5) inhibitor, primarily utilized to improve erectile function. Dapoxetine (60 mg): A selective serotonin reuptake inhibitor (SSRI), employed to help manage premature ejaculation (PE) by increasing control over ejaculation timing. Mechanism of Action Tadalafil Component: This ingredient targets the PDE5 enzyme. By inhibiting PDE5, tadalafil promotes the accumulation of cGMP in the penis during sexual arousal, leading to relaxation of arterial smooth muscles and enhanced blood circulation necessary for achieving and maintaining an erection. Its effects can provide a window of readiness for up to 36 hours. Dapoxetine Component: This ingredient works by altering the levels of serotonin, a neurotransmitter involved in ejaculation control, within the nervous system. By inhibiting serotonin reuptake, dapoxetine helps delay the ejaculatory reflex, offering improved duration before climax. Dapoxetine's effects typically begin within 1-2 hours and are shorter-lasting than tadalafil's. Brand vs. Generic Status Super Vidalista is categorized as a branded generic combination drug. It incorporates active ingredients found in separate established medications: tadalafil is the active component of Cialis® (for ED), and dapoxetine is the active component of Priligy® (for PE, though not approved in all regions like the US). There is no single originator brand that combines these specific two active ingredients in one tablet. Available Forms The medication is supplied as oral tablets, designed for swallowing. Each tablet contains the fixed-dose combination of 20 mg tadalafil and 60 mg dapoxetine. Tablet appearance (color, shape, markings) may vary by manufacturer. Unique Features Dual-Purpose Formulation: Its primary advantage is addressing both erectile dysfunction and premature ejaculation within a single medication dose. Synergistic Approach: Combines tadalafil's long-acting erection support with dapoxetine's targeted action on ejaculation control. Treatment Simplification: Offers convenience by potentially reducing the need for taking separate pills for ED and PE. As-Needed Dosing: Designed for situational use before anticipated sexual activity, rather than daily intake.

---

## General Instructions

**How to Take Super Vidalista Administration:** Ingest one Super Vidalista tablet orally, accompanied by a substantial glass of water. Consume the tablet intact; do not attempt to split, crush, or chew it, as this can alter how the active ingredients are released. **Food Considerations:** It is advisable to avoid consuming large or particularly high-fat meals shortly before taking Super Vidalista. Such meals might impede the absorption of tadalafil and potentially delay the onset of its effects. **Timing Guidelines Optimal Timing:** For optimal efficacy in addressing both ED and PE, take the tablet approximately 1 to 3 hours before planning sexual engagement. This timeframe allows sufficient time for both tadalafil and, more critically, dapoxetine to reach effective levels. **Duration of Effect:** While tadalafil's contribution to erection readiness can persist for up to 36 hours, the effect of dapoxetine on ejaculation control is shorter-lived and more closely tied to this 1-3 hour window pre-activity. **Frequency Maximum Dosage:** Under no circumstances should you exceed one Super Vidalista tablet within a single 24-hour period. **Missed Dose Situational Use:** Super Vidalista is intended for 'on-demand' use prior to sexual relations, not as a regularly scheduled medication. If you plan to use it and miss the intended time, simply skip that dose. Take the next dose when needed, ensuring at least 24 hours have elapsed since the last administration. **Storage Guidelines Conditions:** Maintain the tablets in a controlled room temperature setting, ideally remaining below 30°C (86°F). **Protection:** Preserve the tablets within their original packaging (blister pack) until administration. This safeguards them from environmental factors like moisture and light. Keep the medication securely stored where children cannot access it.

---

## Side Effects

Taking Super Vidalista involves exposure to two active ingredients, tadalafil and dapoxetine, each carrying its own profile of potential side effects. Some individuals might experience adverse reactions, although many tolerate the combination well. **Common Side Effects:** These effects are reported more frequently and are often mild to moderate, sometimes lessening with adaptation to the medication. **Nausea:** A feeling of wanting to vomit, often associated with the dapoxetine component. **Headache:** A common occurrence linked to both tadalafil (due to blood flow changes) and dapoxetine. Staying hydrated might help. **Dizziness:** Feeling lightheaded or unsteady, particularly noted with dapoxetine. Rise slowly from sitting or lying positions. **Facial Flushing:** Temporary skin redness or warmth, primarily due to tadalafil's vasodilating effect. **Diarrhea:** Loose stools may occur, sometimes related to dapoxetine. **Less Common Side Effects:** These reactions are observed less often but can still occur. **Indigestion/Stomach Upset:** Discomfort in the upper abdomen, often linked to tadalafil. **Fatigue/Tiredness:** Feeling unusually weary or lacking energy, potentially from dapoxetine. **Insomnia:** Difficulty falling asleep or staying asleep. **Back or Muscle Soreness:** Aches associated with tadalafil, typically appearing hours later and resolving within a day or two. **Nasal Congestion:** A stuffy nose sensation from tadalafil's effect on blood vessels. **Blurred Vision:** Minor, temporary changes in eyesight. **Serious Side Effects:** These are infrequent but necessitate immediate medical evaluation. **Cease taking Super Vidalista and seek urgent help if you encounter:** **Priapism (Prolonged Erection):** An erection lasting over 4 hours, painful or not. This requires emergency treatment to prevent permanent penile damage (related to tadalafil). **Sudden Vision Loss:** Abrupt decrease or loss of sight in one or both eyes (potential NAION, linked to tadalafil). **Sudden Hearing Impairment:** Rapid decline in hearing, possibly with ringing (tinnitus) or vertigo (associated with tadalafil). **Fainting (Syncope) or Severe Dizziness:** Particularly upon standing (orthostatic hypotension), potentially heightened by the combination or dapoxetine. **Serotonin Syndrome:** A rare but dangerous condition linked to dapoxetine (especially if combined with other serotonergic drugs), with symptoms like agitation, confusion, rapid heartbeat, high fever, muscle rigidity, tremors, or seizures. **Severe Allergic Reactions:** Signs including skin rash, hives, intense itching, difficulty breathing, or significant swelling of the face, lips, tongue, or throat. **Mood Changes:** Significant shifts in mood, depression, or suicidal thoughts (rarely associated with SSRIs like dapoxetine). **Management Note:** If any side effect proves particularly bothersome, persists unexpectedly, or causes significant concern, consult your healthcare provider for advice. Do not alter the dosage or frequency without professional guidance.

---

## Uses

Approved Medical Applications Super Vidalista is specifically indicated for adult men who experience both erectile dysfunction (ED) and premature ejaculation (PE) concurrently. It aims to provide a singular treatment approach for individuals facing this dual diagnosis. How It Addresses These Conditions Combating Erection Difficulties: The tadalafil component functions by inhibiting the PDE5 enzyme, which helps relax penile blood vessels upon sexual stimulation. This improved vascular response allows for enhanced blood inflow, crucial for achieving and sustaining a firmer erection suitable for intercourse over an extended potential timeframe (up to 36 hours). Improving Ejaculatory Control: The dapoxetine component operates as a selective serotonin reuptake inhibitor (SSRI). By modulating serotonin activity in the nervous system pathways controlling ejaculation, it helps to prolong the time before climax occurs, offering better control during sexual activity. Its effect is more time-sensitive, typically peaking within 1-3 hours after intake. Integrated Therapeutic Solution: Super Vidalista offers the convenience of addressing two separate conditions with a single tablet, potentially simplifying the treatment regimen for men diagnosed with both ED and PE. Complementary Action Timing: The formulation combines the long-lasting potential of tadalafil for erection readiness with the relatively rapid-acting effect of dapoxetine targeted at improving ejaculatory latency during the sexual encounter. Important Considerations Symptom Relief, Not Resolution: It's essential to understand that Super Vidalista manages the symptoms of ED and PE but does not cure the underlying physiological, psychological, or relational causes contributing to these conditions. Requires Sexual Arousal: Tadalafil's effect on erections is dependent on the presence of sexual stimulation; the medication does not spontaneously induce erections. Dual Diagnosis Necessary: This combination product is intended only for men formally diagnosed with both erectile dysfunction and premature ejaculation. It is not appropriate for men experiencing only one of these conditions. Comprehensive Medical Evaluation: Use of Super Vidalista should only follow a thorough assessment by a healthcare professional to confirm the dual diagnosis and evaluate suitability, considering potential interactions and individual health status. Not for Recreational Use: This medication should not be used for recreational purposes or by individuals not experiencing both ED and PE.

---

## Safety advice

### Interactions Alcohol:

- Use with caution
- Consuming alcohol while taking Super Vidalista can potentiate side effects associated with both ingredients, such as increased dizziness, headache, rapid heartbeat, or drops in blood pressure (particularly orthostatic hypotension linked to dapoxetine). Significant alcohol intake should be avoided to minimize these risks and ensure medication effectiveness.

### Interactions Other Medications:

- Consult your doctor
- Super Vidalista has a complex interaction profile due to its two active components. It is absolutely contraindicated with nitrates (due to tadalafil). Additionally, significant interactions exist with other PDE5 inhibitors, strong CYP3A4 inhibitors/inducers (affecting tadalafil), alpha-blockers, antihypertensives, and crucially, other serotonergic drugs (SSRIs, SNRIs, MAOIs, triptans, St. John's Wort) which increase the risk of serotonin syndrome when combined with dapoxetine. A complete medication review by a physician is mandatory.

### Special Groups Pregnancy:

- Unsafe
- Super Vidalista is formulated strictly for use in adult men dealing with ED and PE. It is absolutely not intended for, nor should it be used by, women, especially those who are pregnant.

### Special Groups Breastfeeding:

- Unsafe
- This combination medication is contraindicated for women, including those who are breastfeeding. Its use is confined to managing specific male sexual health conditions.

### **Special Groups Elderly:**

- Use with caution
- Older males (typically 65+) may be more susceptible to the side effects of both tadalafil (e.g., blood pressure effects) and dapoxetine (e.g., dizziness, orthostatic hypotension). Careful medical consideration of overall health, kidney function, and potential for falls is necessary before prescribing Super Vidalista to this age group.

### **Special Groups Children:**

- Unsafe
- Super Vidalista is strictly prohibited for use in individuals under 18 years of age. Its safety and efficacy have not been established in pediatric or adolescent populations.

### **Effects on Activities Driving:**

- Use with caution
- The potential for dizziness, blurred vision (from tadalafil), fatigue, or fainting (syncope, particularly linked to dapoxetine) necessitates caution when driving. Assess your individual reaction thoroughly before operating a vehicle after taking Super Vidalista.

### **Effects on Activities Operating Machinery:**

- Use with caution
- Similar to driving, activities demanding high levels of alertness and coordination, such as operating heavy machinery, should be avoided until you are certain that Super Vidalista does not impair your ability to perform these tasks safely due to potential side effects like dizziness, fatigue, or visual changes.

---

## **Concerns**

**Key Concerns When Considering Super Vidalista Cardiovascular Stress and Hypotension:** The tadalafil component can lower blood pressure, while dapoxetine can contribute to orthostatic hypotension (dizziness/fainting upon standing). This combination poses risks for individuals with certain cardiovascular conditions, those taking antihypertensives, or those prone to fainting. Sexual activity itself adds strain; hence, careful cardiovascular evaluation is needed. **Risk of Serotonin Syndrome:** Dapoxetine, being an SSRI, increases serotonin levels. Combining Super Vidalista with other substances that also boost serotonin (like other antidepressants, certain pain medications, St. John's Wort, or illicit drugs) significantly elevates the risk of serotonin syndrome – a potentially life-threatening condition characterized by agitation, fever, rapid heartbeat, and muscle rigidity. **Priapism Potential:** Tadalafil carries the rare but serious risk of causing priapism (an erection lasting over 4 hours). This requires emergency medical treatment to prevent long-term damage to the penis. **Neuropsychiatric Effects:** Dapoxetine, like other SSRIs, has been associated with potential neuropsychiatric effects, including dizziness, fatigue, insomnia, and rarely, mood changes or fainting (syncope). Individuals with a history of psychiatric conditions or fainting spells require careful assessment. **Impact of Impaired Organ Function:** Both tadalafil and dapoxetine are metabolized by the liver and excreted via the kidneys. Significant impairment in either organ's function can lead to increased drug levels and heightened risk of adverse effects. Dose adjustments or avoidance may be necessary. **When Not to Use Super Vidalista Absolute Contraindications:** Use is strictly forbidden with nitrates (due to tadalafil). It's also contraindicated in individuals with significant heart pathology (e.g., heart failure, conduction abnormalities, significant valve disease), a history of fainting (syncope), mania or severe depression, known hypersensitivity to either tadalafil or dapoxetine, and severe liver impairment. Concurrent use with potent CYP3A4 inhibitors (affecting tadalafil) or MAO inhibitors (affecting dapoxetine) is also prohibited. Use in severe kidney disease is generally not recommended. **Critical Reminders Dual Diagnosis**

**Prerequisite:** Emphasize that Super Vidalista is exclusively intended for men diagnosed with both erectile dysfunction and premature ejaculation by a qualified healthcare provider. It is not for treating either condition alone. **Beware Counterfeit Combination Drugs:** Acquiring combination medications like Super Vidalista from unverified sources poses a significant danger. Fakes might contain incorrect ratios of ingredients, substitute substances, or harmful contaminants, drastically increasing health risks. Rely only on legitimate prescriptions and suppliers. **Avoid Abrupt Discontinuation (if used regularly):** While typically used ‘as needed’, if taken more regularly under specific medical advice (uncommon), stopping SSRIs like dapoxetine abruptly can sometimes lead to withdrawal symptoms. Follow medical guidance on cessation. **Report Serious Symptoms Promptly:** Do not delay seeking medical attention for warning signs like fainting, severe mood changes, symptoms suggestive of serotonin syndrome, priapism, or sudden vision/hearing loss.

---

## Warnings

**Critical Health Authority Precautions** Regulatory agencies highlight crucial safety aspects for the components of Super Vidalista: **Tadalafil – Nitrate Hazard:** A primary warning emphasizes the severe risk of life-threatening hypotension if tadalafil is combined with any organic nitrates. **Tadalafil – Cardiovascular Assessment:** Evaluation of heart health is essential before prescribing tadalafil due to its vascular effects. **Tadalafil – Sudden Vision Loss Risk (NAION):** Rare occurrences of non-arteritic anterior ischemic optic neuropathy leading to vision impairment are linked to PDE5 inhibitors. **Dapoxetine – Serotonin Syndrome Risk:** Co-administration with other serotonergic agents significantly increases the risk of potentially fatal serotonin syndrome. **Dapoxetine – Neuropsychiatric Risks:** Potential for dizziness, fainting (syncope), and mood changes requires careful patient selection and monitoring. (Note: Dapoxetine itself is not FDA-approved in the US). **Absolute Contraindications** Super Vidalista must never be used under the following conditions: **Concurrent Nitrate Use:** Absolutely forbidden alongside any form of nitrate medication (e.g., nitroglycerin) or recreational “poppers.” **MAOI Use:** Taking Super Vidalista concurrently with, or within 14 days of stopping, Monoamine Oxidase Inhibitors (MAOIs) is prohibited due to the risk of severe reactions including serotonin syndrome. **Potent CYP3A4 Inhibitor Use:** Co-administration with strong inhibitors like ketoconazole, itraconazole, or ritonavir is contraindicated as it significantly increases tadalafil levels. **Significant Cardiac Pathology:** Patients with certain types of heart failure, heart block or sick sinus syndrome (unless a permanent pacemaker is in place), significant valvular disease, or a history of unstable angina/recent major cardiac event should not use this drug. **History of Syncope, Mania, or Severe Depression:** Individuals with a documented history of fainting spells, manic episodes, or severe depressive illness are contraindicated. **Known Hypersensitivity:** Prior serious allergic reaction to either tadalafil or dapoxetine, or any tablet excipients. **Severe Liver Impairment:** Not suitable for individuals with severely compromised liver function. **Urgent Precautions** Immediate medical attention is required if the following develop after taking Super Vidalista: **Priapism:** An erection persisting for more than 4 hours warrants emergency treatment. **Sudden Sensory Loss:** Abrupt loss or reduction of vision or hearing requires immediate cessation and medical care. **Fainting (Syncope):** Any episode of losing consciousness after taking the medication needs urgent evaluation. **Symptoms Suggesting Serotonin Syndrome:** Development of agitation, confusion, high fever, rapid heartbeat, sweating, tremors, or muscle stiffness demands emergency medical assessment. **Severe Cardiovascular Symptoms:** Experiencing chest pain, severe dizziness, or irregular heartbeat during or after sexual activity requires immediate medical help.

---

## Dosage

**Standard Dosing Guidelines** **Recommended Dose:** The standard dose is one Super Vidalista tablet, containing tadalafil 20 mg and dapoxetine 60 mg, taken orally as required. **Administration:** Swallow the tablet whole with a full glass of water. Do not break, chew, or crush it. **Timing:** Ingest the tablet approximately 1 to 3 hours prior to planned sexual activity to allow both components to become effective. **Maximum Frequency:** Strictly limit intake to no more than one tablet within any 24-hour duration. **Dose Adjustments** **Renal Impairment:** For mild or moderate kidney problems, Super Vidalista should be used with caution. Its

use is generally not recommended in individuals with severe kidney impairment (e.g., estimated Glomerular Filtration Rate [eGFR] < 30 mL/min) due to limited data and clearance concerns for both active ingredients. Always follow medical advice. Hepatic Impairment: Use Super Vidalista with caution in cases of mild liver dysfunction. However, due to the dapoxetine component, this medication is contraindicated (must not be used) in individuals suffering from moderate or severe hepatic impairment. Elderly Patients: Older adults (typically 65+) may experience increased sensitivity to side effects from both tadalafil (like blood pressure changes) and dapoxetine (like dizziness or fainting). Super Vidalista should be prescribed cautiously to elderly patients only after careful medical evaluation of potential risks versus benefits. Missed Dose Super Vidalista is intended for 'on-demand' usage before sexual intimacy, not regular daily administration. If you forget to take a planned dose, simply skip it. Take the next dose when needed, provided it's been at least 24 hours since the last one. Overdose Management Symptoms: Taking excessive amounts of Super Vidalista can lead to intensified side effects from both components. Potential signs include severe headache, prolonged erection (priapism), significant dizziness or fainting, pronounced nausea or vomiting, rapid heartbeat, tremors, agitation, or symptoms possibly indicative of serotonin syndrome. Action: A suspected overdose requires immediate emergency medical attention. Contact emergency services or a poison control center without hesitation. Treatment will focus on supportive care based on the observed symptoms.

---

## Interactions

**Drug-Drug Interactions** Nitrates (Hazardous Prohibition): Co-use of Super Vidalista with any organic nitrate medication (e.g., nitroglycerin, isosorbide compounds used for angina) or recreational nitrites ("poppers") is absolutely contraindicated. This combination poses a severe risk of precipitous, life-threatening hypotension due to tadalafil. MAO Inhibitors (MAOIs): Combining Super Vidalista with MAOIs (or using within 14 days of stopping an MAOI) is strictly forbidden due to the dapoxetine component, risking serious, potentially fatal reactions including serotonin syndrome. Similarly, MAOIs should not be started within 7 days of stopping Super Vidalista. Other Serotonergic Agents: Extreme caution is necessary, or combinations should be avoided, with other drugs that increase serotonin levels (e.g., other SSRIs, SNRIs, triptans for migraines, tramadol, linezolid, lithium) due to the elevated danger of serotonin syndrome caused by dapoxetine. Alpha-Blockers & Antihypertensives: Using Super Vidalista alongside alpha-blockers (for BPH/hypertension) or other blood pressure-lowering medicines can lead to additive hypotensive effects, increasing the chance of dizziness or fainting. Careful medical management is required. Potent CYP3A4 Inhibitors: Strong inhibitors of the CYP3A4 enzyme (like ketoconazole, itraconazole, ritonavir) can significantly boost levels of tadalafil and potentially dapoxetine. Co-administration with potent CYP3A4 inhibitors is generally contraindicated or requires significant caution/dose adjustment as per medical guidance. Thioridazine: Administration of thioridazine is contraindicated with dapoxetine due to the risk of serious heart rhythm problems (QTc prolongation). Wait appropriate intervals when switching between these drugs. Other PDE5 Inhibitors / Riociguat: Concurrent use with other ED treatments (sildenafil, vardenafil) or riociguat is not advisable due to cumulative side effects, particularly hypotension.

**Drug-Food Interactions** Substantial High-Fat Meals: Consuming a meal very high in fat content just before taking Super Vidalista might potentially delay the absorption and onset time, particularly for the tadalafil component. Grapefruit / Grapefruit Juice: Significant intake of grapefruit products may inhibit CYP3A4 metabolism, potentially increasing exposure to tadalafil and possibly dapoxetine, thus raising side effect risks. Moderation or avoidance is often suggested. Alcohol: Combining alcohol with Super Vidalista is discouraged. Alcohol can worsen orthostatic hypotension (dizziness/fainting on standing, associated with dapoxetine), intensify dizziness or drowsiness, and independently impair both erectile function and judgment.

**Drug-Supplement/Herb Interactions** St. John's Wort (*Hypericum perforatum*): This herbal supplement can induce CYP3A4 (potentially reducing tadalafil effectiveness) and also possesses serotonergic properties, increasing the risk of serotonin syndrome when taken with dapoxetine. Concurrent use should be avoided. Supplements Affecting Serotonin or Blood Pressure: Caution is advised with supplements known to affect serotonin levels (e.g., tryptophan, 5-HTP) or blood pressure, as they might interact additively with Super Vidalista's components. Impact of Interactions Interactions involving Super Vidalista can range from reduced treatment effectiveness to significantly heightened risks of adverse events like severe low blood pressure, serotonin

syndrome, increased component-specific side effects, or fainting. Some interactions represent immediate medical emergencies. **Critical Reminders Mandatory Full Disclosure:** To ensure safety, it is imperative to provide your healthcare provider with a totally comprehensive inventory of all substances used—including prescription/non-prescription drugs, herbal remedies, supplements, and recreational drugs—before starting Super Vidalista. **Counterfeit Combination Interaction Hazards:** Be acutely aware that illicitly sourced Super Vidalista may contain incorrect dosages or ratios of tadalafil and dapoxetine, or even entirely different undeclared substances, creating unpredictable and potentially severe interaction risks with your other medications.

---

## FAQs

- **Can Super Vidalista’s dual composition (tadalafil + dapoxetine) cause stomach upset?**  
Yes, Super Vidalista’s combination of tadalafil (for erectile dysfunction) and dapoxetine (for premature ejaculation) may cause gastrointestinal issues like nausea, indigestion, or diarrhea in some users. These side effects are typically mild but can vary depending on individual sensitivity.
  - **Is Super Vidalista safe when combined with acupuncture?**  
Super Vidalista itself doesn’t directly interact with acupuncture. However, always inform your acupuncturist about medications you’re taking. For example, if acupuncture targets stress or blood flow (common in sexual health therapies), ensure your practitioner knows about Super Vidalista to avoid overlapping effects. Safety first!
  - **Does Super Vidalista worsen autoimmune disease symptoms (e.g., arthritis, lupus)?**  
There’s limited research on Super Vidalista’s impact on autoimmune conditions. However, tadalafil (a PDE5 inhibitor) may affect blood flow and inflammation pathways, which could theoretically interact with autoimmune responses.
  - **Can long-term Super Vidalista use lead to emotional dependency?**  
Super Vidalista isn’t addictive, but relying on it for confidence in intimate situations might create psychological dependence over time. Some users report feeling anxious about performance without the medication. If you notice emotional changes, consider counseling or therapy alongside medical treatment to address underlying concerns.
- 

## Other Details

**Appearance** Super Vidalista tablets, containing 20 mg tadalafil and 60 mg dapoxetine, are typically oral film-coated tablets. Their specific color, shape (e.g., oval, round), and any score lines or embossed markings can differ depending on the manufacturing source (verify packaging details; Centurion Laboratories is often associated with this product). **Storage After Opening** It is recommended to keep Super Vidalista tablets securely within their original blister packaging until the intended time of use. This practice helps maintain their stability by protecting them from environmental factors like humidity and light exposure. Avoid transferring tablets to multi-dose pill organizers. **Disposal** Expired or no longer needed Super Vidalista tablets should not be disposed of via flushing down drains or toilets. Investigate local drug take-back schemes offered by pharmacies if available. If not, render the tablets undesirable (e.g., mix with an unpalatable substance like coffee grounds), seal them within a plastic bag, and discard with regular household waste. **Lab Test Interference** Tadalafil’s known effect is lowering blood pressure, which might affect readings during medical checks; inform healthcare staff accordingly. Dapoxetine does not typically cause significant interference with common lab tests, but always disclose all medication use to laboratory personnel and physicians. **Handling Precautions** Take the tablet whole with water as directed. Given the combination of active ingredients with specific indications and potential side effects, it is particularly important to ensure this medication is stored securely away from children, pets, and women (especially pregnant or breastfeeding individuals). **Counterfeit Identification Tips** Packaging Examination: Authentic

medication usually features high-quality packaging with clear printing, verifiable batch numbers, expiry dates, and possibly security features (holograms, seals). Be suspicious of poorly printed boxes, damaged blister packs, or missing information. **Tablet Uniformity:** Genuine tablets should exhibit consistency in color, shape, size, and coating. Variations, unusual textures (chalky, crumbly), or inconsistent markings could indicate a counterfeit product. **Source Validation:** The risk of acquiring fake or substandard combination drugs like Super Vidalista is heightened when purchasing from unregulated online vendors or sources outside legitimate pharmacy channels. Always use a prescription and obtain medication from trusted suppliers. **Additional Advice** If traveling, carrying Super Vidalista in its original packaging with a corresponding prescription copy can prevent potential issues with customs or authorities. Be mindful of the expiration date; using expired medication may result in reduced efficacy or potential safety concerns. Report any perceived quality issues with the tablets (e.g., unusual smell, appearance) to your pharmacist before consumption. Other formulations are available within the Vidalista product line to suit different patient requirements, including: Vidalista Black Vidalista Professional Vidalista CT

---

## References

Drugs.com: Tadalafil. Drugs.com; 2024. [Updated Apr 22, 2024]. [Accessed May 21, 2024]. <https://www.drugs.com/tadalafil.html> MedlinePlus: Tadalafil. U.S. National Library of Medicine; 2024. [Updated Apr 15, 2024]. [Accessed May 21, 2024]. <https://medlineplus.gov/druginfo/meds/a604008.html> Mayo Clinic: Tadalafil (Oral Route) Description. Mayo Foundation for Medical Education and Research; 2024. [Updated Feb 1, 2024]. [Accessed May 21, 2024]. <https://www.mayoclinic.org/drugs-supplements/tadalafil-oral-route/description/drg-20067204> WebMD: Tadalafil Oral: Uses, Side Effects, Interactions, Pictures, Warnings & Dosing. WebMD; 2024. [Accessed May 21, 2024]. <https://www.webmd.com/drugs/2/drug-77875/tadalafil-oral/details>

---

## Disclaimer

Disclaimer: The information on this site is provided for informational purposes only and is not medical advice. It does not replace professional medical consultation, diagnosis, or treatment. Do not self-medicate based on the information presented on this site. Always consult with a doctor or other qualified healthcare professional before making any decisions about your health.