

# iMedix: Your Personal Health Advisor.

## Tadalista® (Tadalafil)

---

Tadalista is aimed at increasing blood flow to the penis resulted from the relaxation of smooth muscles in it. The main active ingredient is Tadalafil.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

### Description

General information Tadalista is a generic drug whose branded equivalent is Cialis. The manufacturer of the medication is the company Fortune Health Care, which deals in various superior quality generic medicines from top brands for treating loss of sexual power. Tadalista is among the most widely known drugs being employed today for improving men's health. It is qualified as a miracle because it starts working as fast as in 30 min after dosing with an outcome that can go on for 36 hours. A man who uses this drug will not need to be anxious about going on a weekend journey in company with his lover merely because he will be able to retain a stiff male organ for almost 2 days. Tadalista's main component is Tadalafil, a chemical that is categorized as PDE-5. This ingredient acts by helping the blood flow to reach the male organ and therefore prolongs the hard-on. The stimulant action of this remedy is more efficacious than in other similar pharmaceuticals, like Filitra and Filagra (such as sildenafil, tadalafil, vardenafil, kamagra, suhagra, tadacip, revatio, cenforce). Unlike other loss of sexual power drugs, which action lasts only for six hours, Tadalista's motivational effect helps to feel confident for 36 hours. Precautions During intercourse, the load on the cardiovascular system increases significantly. This can be fatal if there are serious heart health problems. Taking many medications (including Tadalista) can multiply the demonstration of adverse reactions and carry an additional load on the heart at the time of sex. One must urgently seek medical aid if the following repercussions occur: severe vertiginousness; collapse; nausea; numbness of upper and lower limbs; stethalgia or idiopathic hypertension; feeling short of breath; left-handed pain; sudden deterioration of vision in one or both eyes (including blindness). Impairments of vision are more likely to occur to those who suffer from pancreatic diabetes, circulation diseases, have elevated levels of cholesterol in the blood, idiopathic hypertension, are older than 60 years, as well as men who have a cigarette from time to time. A sudden decrease in hearing or diminished hearing, tinnitus, and giddiness are observed. In this case, it is recommended to stop using Tadalista and seek medical assistance immediately. On the background of taking the remedy, the onset of a long (above 4-5 hours) and painful erection is eventual. If after that you do not give up administering the medication, you may experience serious health problems afterward. Severe inordinate responses to this medicament are rare, but inactive ingredients that makeup it can cause them. Therefore, in the event of exanthematous disease and skin irritation, tumour of the face, neck, tongue, shortness of breath, medical care is required immediately. Before using Tadalista, you should inform your health adviser about acute or chronic diseases that you have, such as heart conditions (for example, an acute coronary syndrome or stroke, life-threatening arrhythmias, chest pain/ coronarism, cardiac impairment), high

or low blood tension, fibrosis or scarring, plastic induration, a painful or too long erection (priapism) experienced in the past, leucosis or multiple myeloma, eye diseases (primary pigmentary degeneration of retina, vision impairment), disorders of the blood clotting system in an active phase of gastric ulcer. Possible giddiness when taking Tadalista is a contraindication to driving a car and mechanisms that require increased concentration of attention. When taking the drug, one should limit the intake of alcoholic beverages. If it became necessary to perform surgically (also dental) operations, you must tell the medical adviser about all the medicinal and herbescent remedies that you use.

**What is Tadalista?** Tadalista is a medication used for treating erectile dysfunction (ED) in men. It contains tadalafil as its active ingredient, which helps in achieving and maintaining an erection by improving blood flow to the penis.

**How does Tadalista work?** Tadalista works by inhibiting the phosphodiesterase type 5 (PDE5) enzyme, leading to increased blood flow to the penis during sexual arousal. This helps to achieve and maintain an erection suitable for sexual activity.

**What are the available dosages of Tadalista?** Tadalista is available in various dosages, including Tadalista 10 mg, 20 mg, 40 mg, and Tadalista Super Active. The appropriate dosage depends on the individual's health condition and the severity of their ED.

**What are the possible side effects of Tadalista?** Common side effects of Tadalista include headache, back pain, muscle aches, nasal congestion, indigestion, and facial flushing. Rare but serious side effects can include changes in vision, sudden hearing loss, and an erection lasting more than four hours (priapism).

**Can Tadalista be taken with alcohol?** It is advisable to limit or avoid alcohol while taking Tadalista, as alcohol can increase the risk of side effects such as dizziness and can also impair sexual performance.

**How should Tadalista be taken?** Tadalista should be taken orally, with or without food, about 30 minutes before sexual activity. The effects of Tadalista can last for up to 36 hours. Only one tablet should be taken per day.

**Is Tadalista suitable for everyone?** Tadalista is not suitable for everyone. Men with certain health conditions like heart disease, low blood pressure, or those taking nitrates should not take Tadalista. Always consult with a healthcare provider before starting this medication.

**Can Tadalista cure erectile dysfunction permanently?** Tadalista is not a cure for erectile dysfunction. It is a treatment that helps to manage the symptoms of ED by enabling the achievement of an erection for sexual activity.

**Does Tadalista require a prescription?** Yes, Tadalista is a prescription medication. It is important to consult a healthcare provider to get a prescription and to ensure that Tadalista is an appropriate treatment option.

**Can Tadalista interact with other medications?** Tadalista can interact with certain medications, especially nitrates used for chest pain, alpha-blockers, and some antifungal drugs. Always inform your healthcare provider about all medications and supplements you are taking.

**Dosage** The remedy comes in 3 variants: 20, 10, and 5 mg. The doses define the efficacy comparable with each pill size. Apart from the pill, there are also Tadalista chewy pill, famous for its continuous exposure of over a day. The recommended daily intake is 20mg. The measuring hinges on the patient's state of health, the body's response to treatment, and reciprocal interaction with other pharmaceuticals currently being taken. Before the appointment, it is important to inform the physician about all the drugs used, including those of plant origin, dietary supplements, vitamins.

In the complex therapy of Prostatic Hyperplasia, the medicine is taken according to the prescribed scheme, usually once every 24 hours. If Finasteride is also taken in the therapy complex, the duration of its administration should be determined by the attending physician. For the treatment of sexual debility by Tadalista, two methods are used, the choice of which is selected individually. The first way is to take the remedy as needed, as a rule, at least 30-40 min before sexual intercourse. As it was said already, Tadalista exposure to a man's sexual abilities may last for about 36 hours. With another treatment for loss of sexual power, Tadalista is taken on a regular basis every day, once every 24 hours. If the drug is taken regularly, the opportunity to make sexual contact can be exercised at any time between the administration of the pill. In the case of simultaneous treatment of invirility and benign prostatic hyperplasia, Tadalista is usually taken once a day. In this sexual intercourse can be carried out by a man at the desired time between taking the next dose of the drug. It is important not to miss the next dose. How to take Tadalista should be taken with a glass of water or other non-alcoholic liquid. If the remedy is taken on a daily basis and the next dose has been missed, you must take it as soon as possible. If this moment coincided with the time of the next dose, then you should not take the skipped pill, doubling it, after that you need to recur to the usual regimen.

**Side-effects** Despite the fact that the effect of Tadalista is quite mild, and the drug is tested and completely safe, the feasibility of undesirable sequences during the reception of the drug cannot be ruled out. In particular, in rare cases, there may occur: A headache and muscle aches; A rhinorrhea; Vertiginousness; Flushing of the face; Tightening of lower and upper limbs; Hypertonia. In the words of the health consumers, adverse effects from taking the

remedy are extremely rare, while the unpleasant indicants vanish shortly after stopping the drug and do not need special treatment. Overdose In the case of overdose, the same side-effects are most often manifested as with lower doses: a headache, dizziness, fainting, feeling short of breath, etc. If an overdose is suspected, it is necessary to contact a first-aid station. Interactions If you conceive an idea to buy and begin taking Tadalista capsules, remember that there are a number of restrictions for taking them. So, Tadalafil containing drugs cannot be used: by patients under 18; the first six months after a stroke or heart seizure; by people with renal or hepatic failure; in case of intolerance or hypersensitivity to the components of the product; in case of acute heart disease. Tadalista can cause a significant reduction in blood tension while taking nitrates (such as molsidomine, amyl nitrite). This can cause severe dizziness, fainting, and sometimes a heart attack or stroke. Tadalista should not be taken with medicines used to treat chest pain/stenocardia, or recreational medicines containing amyl or butyl nitrite. If Tadalista is taken along with alpha-blocking drugs (such as doxazosin, tamsulosin) to treat high blood pressure, an enlarged prostate, or prostate adenoma, blood pressure can drop significantly, which can lead to fainting and severe giddiness. In this case, treatment with a lower dose of Tadalista is prescribed or the dose of alpha-blocking drugs is adjusted in order to minimize the risk of a drop in blood pressure. Some pharmaceuticals may affect the efficacy and speed of excretion of Tadalista from the body. These include azidimidazole (itraconazole, ketoconazole), macrolides (clarithromycin, erythromycin), HIV protease inhibitors (fosamprenavir, ritonavir), hepatitis virus with protease inhibitors (boceprevir, telaprevir), rifampin, and others. You should not take the medicaments that include active substance tadalafil, or other means to treat erectile dysfunction or lesser circuit hypertension (such as sildenafil, vardenafil). Alcohol interaction Despite the fact that Tadalista can be combined with taking small doses of alcohol, it is still better to refuse alcohol because it increases the risk of side-effects. Reviews Boris (51): I noticed slight dizziness after application. The effect is good, but after its occurrence, you need to sit or lie down for 15-20 minutes so that everything will return to normal. Perhaps the tool does not suit me personally. However, the main task is coping just fine; my woman is more satisfied than ever. Klaus (37): Tadalista is generally an effective medicine, particularly if you drink it on an empty stomach, after thirty minutes I was fully prepared. I heard that there are often fakes, be more careful. Samuel Lee (44): I tried Tadalista for the first time, after reading about its magical properties on the Internet. Also, there were very good reviews of this drug. However, I didn't notice any effect. The erection is the same as it was before. I caught a side effect, on the second day I could not fall asleep from pain in my chest. I was forced to call an ambulance. Therefore, I would not recommend Tadalista to anyone. John (54): I bought and took Tadalista 20 mg a couple of days ago. If you order this pill, get ready for the fact that its action is powerful. However, I had side-effects, such as a headache and dizziness.

---

---

## Side Effects

---

## Uses

---

---

---

---

---

## **Interactions**

---

---

## **Other Details**

---

---