

iMedix: Your Personal Health Advisor.

Talquetamab-Tgvs 2 Mg/Ml Subcutaneous Solution

Search for medical details concerning talquetamab-tgvs subcutaneous on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Tiredness, nail problems, muscle/joint pain, loss of appetite, weight loss, dry mouth/eyes/skin, or diarrhea may occur. Pain, bruising, redness, itching, or swelling at the injection site may also occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Pain or sores in the mouth and throat may occur. Brush your teeth carefully/gently, avoid using mouthwash that contains alcohol, and rinse your mouth often with cool water mixed with baking soda or salt. It may also be best to eat soft, moist foods. People using this medication may have serious side effects. However, you have been prescribed this drug because your doctor has judged that the benefit to you is greater than the risk of side effects. Careful monitoring by your doctor may decrease your risk. Tell your doctor right away if you have any serious side effects, including: easy bruising/bleeding, signs of liver disease (such as nausea/vomiting that doesn't stop, stomach/abdominal pain, yellowing eyes/skin, dark urine), signs of anemia (such as unusual tiredness, pale skin), oral problems (changes in sense of taste, trouble swallowing). This medication may lower your ability to fight infections. This may make you more likely to get a serious (rarely fatal) infection or make any infection you have worse. Tell your doctor right away if you have any signs of infection (such as sore throat that doesn't go away, fever, chills, cough). Talquetamab can cause skin problems (such as redness, rash, small bumps) during treatment. However, you may not be able to tell it apart from a rare rash that could be a sign of a severe reaction. Tell your doctor right away if you develop any rash. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. How to use Talquetamab-Tgvs 2 Mg/Ml Subcutaneous Solution Read the Medication Guide provided by your pharmacist before you start using talquetamab and each time you get a treatment. If you have any questions, ask your doctor or pharmacist. Carry the Patient Wallet Card with you at all times. Show the card to all of your health care providers to let them know that you are being treated with talquetamab. This medication is given by injection under the skin as directed by your doctor. The injection is given by a health care professional, usually under the skin of the abdomen or thigh. To reduce your risk of side effects, your doctor will start this medication at a low dose and gradually increase your dose. These starting doses are part

of a “step-up dosing schedule.” You must stay in the hospital for 48 hours after each of the “step-up” doses so your doctor can monitor you closely and help you manage any side effects. The dosage and treatment schedule are based on your medical condition, weight, and response to treatment. Before you receive this medication, your doctor will prescribe other medications (such as acetaminophen, dexamethasone, diphenhydramine) to help prevent side effects. Your doctor may also prescribe other medications (such as antibiotics) to help prevent infection. Carefully follow your doctor’s directions for all your medications.

Precautions Before using talquetamab, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: current/recent infections. This drug may make you dizzy or drowsy or affect your thinking and coordination. Alcohol or marijuana (cannabis) can worsen these effects. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). This medication can make you more likely to get infections or may make current infections worse. Stay away from anyone who has an infection that may easily spread (such as chickenpox, COVID-19, measles, flu). Talk to your doctor if you have been exposed to an infection or for more details. Tell your health care professional that you are using talquetamab before having any immunizations/vaccinations. Avoid contact with people who have recently received live vaccines (such as flu vaccine inhaled through the nose). To lower the chance of getting cut, bruised, or injured, use caution with sharp objects like razors and nail cutters, and avoid activities such as contact sports. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Tell your doctor if you are pregnant or plan to become pregnant. You should not become pregnant while using talquetamab. Talquetamab may harm an unborn baby. Your doctor should order a pregnancy test before you start this medication. Ask about reliable forms of birth control while using this medication and for 3 months after the last dose. If you become pregnant, talk to your doctor right away about the risks and benefits of this medication. It is unknown if this drug passes into breast milk. Because of the possible risk to the infant, breast-feeding while using this drug and for 3 months after the last dose is not recommended. Consult your doctor before breast-feeding. Consult your pharmacist or physician.

Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Lab and/or medical tests (such as complete blood count, liver function, weight) should be done before you start using this medication and while you are using it. Keep all medical and lab appointments. Consult your doctor for more details. It is important to get each dose of this medication as scheduled. If you miss a dose, ask your doctor or pharmacist right away for a new dosing schedule. Not applicable. This medication is given in a hospital or clinic and will not be stored at home.

Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor’s approval.

Side Effects

Uses

Interactions

Other Details
