

iMedix: Your Personal Health Advisor.

Tamsulosin HCL

Search for medical details concerning tamsulosin oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Dizziness, lightheadedness, drowsiness, runny/stuffy nose, or ejaculation problems may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. To reduce the risk of dizziness and lightheadedness, get up slowly when rising from a sitting or lying position. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: fainting. Rarely, males may have a painful or prolonged erection lasting 4 or more hours. If this occurs, stop using this drug and get medical help right away, or permanent problems could occur. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Tamsulosin is used by men to treat the symptoms of an enlarged prostate (benign prostatic hyperplasia-BPH). It does not shrink the prostate, but it works by relaxing the muscles in the prostate and the bladder. This helps to relieve symptoms of BPH such as difficulty in beginning the flow of urine, weak stream, and the need to urinate often or urgently (including during the middle of the night). Tamsulosin belongs to a class of drugs known as alpha blockers. Do not use this medication to treat high blood pressure. How to use Tamsulosin HCL Read the Patient Information Leaflet if available from your pharmacist before you start taking this medication and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth as directed by your doctor, usually once daily, 30 minutes after the same meal each day. Swallow this medication whole. Do not crush, chew, or open the capsules. The dosage is based on your medical condition and response to treatment. Tamsulosin may cause a sudden drop in your blood pressure, which could lead to dizziness or fainting. This risk is higher when you first start taking this drug, after your doctor increases your dose, or if you restart treatment after you stop taking it. During these times, avoid situations where you may be injured if you faint. Take this medication regularly to get the most benefit from it. To help you remember, take it at the same time each day. If you have not taken this drug for several days, contact your doctor to see if you need to be restarted at a lower dose. It may take up to 4 weeks before your symptoms improve. Tell your

doctor if your condition does not improve or if it worsens. **Precautions** Before taking tamsulosin, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: low blood pressure, certain eye problems (cataracts, glaucoma). This drug may make you dizzy or drowsy. Alcohol or marijuana (cannabis) can make you more dizzy or drowsy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery (including cataract/glaucoma eye surgery), tell your doctor or dentist if you are taking or have ever taken this medication, and about all the other products you use (including prescription drugs, nonprescription drugs, and herbal products). Older adults may be more sensitive to the side effects of this drug, especially dizziness and low blood pressure when getting up from a sitting or lying position. These side effects can increase the risk of falling. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. Consult your doctor before breast-feeding. **Overdose** If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: severe dizziness, fainting. Do not share this medication with others. Lab and/or medical tests (such as prostate exams, blood pressure) should be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. **Interactions** Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this drug include: other alpha blocker drugs (such as prazosin, terazosin). If you are also taking a drug to treat erectile dysfunction-ED or pulmonary hypertension (such as sildenafil, tadalafil, vardenafil, kamagra, suhagra, tadalafil, revatio, cenforce), your blood pressure may get too low which can lead to dizziness or fainting. Your doctor may need to adjust your medications to minimize this risk. Other medications can affect the removal of tamsulosin from your body, which may affect how tamsulosin works. Examples include adagrasib, azole antifungals (such as itraconazole, ketoconazole), clarithromycin, cobicistat, HIV protease inhibitors (such as lopinavir), mifepristone, ribociclib, ritonavir, among others.

Side Effects

Uses

Interactions

Other Details
