

iMedix: Your Personal Health Advisor.

Tolnaftate 1 % Topical Cream

Search for medical details concerning tolnaftate topical on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Irritation of the treated skin may occur. If this effect lasts or gets worse, notify your doctor or pharmacist promptly. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** Tolnaftate is used to treat skin infections such as athlete's foot, jock itch, and ringworm. It is an antifungal that works by preventing the growth of fungus. Some forms of tolnaftate may also be used to prevent athlete's foot from coming back. **How to use Tolnaftate 1 % Topical Cream** Use this medication on the skin only. Clean and thoroughly dry the area to be treated. Apply this medication to the affected skin, usually twice a day or as directed on the product package or by your doctor. The dosage and length of treatment depends on the type of infection being treated. Some forms of tolnaftate (such as powder) need to be shaken before applying. Check your product package to see if your form of this medication needs to be shaken. Apply enough medication to cover the affected area and some of the surrounding skin. After applying this medication, wash your hands. Do not wrap, cover, or bandage the area unless directed to do so by your doctor. Do not apply this medication in the eyes, nose, mouth, or vagina. Use this medication regularly in order to get the most benefit from it. Remember to use it at the same time(s) each day. Continue to use this medication until the full prescribed amount is finished, even if symptoms disappear after starting tolnaftate. Stopping the medication too early may allow the fungus to continue to grow, which may result in a return of the infection. Tell your doctor if your condition lasts after 2 weeks of treatment for jock itch, after 4 weeks of treatment for athlete's foot or ringworm, or if your condition gets worse at any time. **Precautions** Before using tolnaftate, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal

products). This medication should be used only if clearly needed during pregnancy. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. This medication has been prescribed for your current condition only. Do not use it later for another infection unless your doctor tells you to. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Side Effects

Uses

Interactions

Other Details
