

iMedix: Your Personal Health Advisor.

Tretinoin 0.025 % Topical Cream

Search for medical details concerning tretinoin topical on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Warmth or stinging may occur right after applying the medication. Skin redness, dryness, itching, scaling, mild burning, or worsening of acne may occur during the first 2 to 4 weeks of using the medication. These effects usually decrease with continued use. A daytime moisturizer may be helpful for very dry skin . If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: blistering/crusting of the skin, severe burning/swelling of the skin, skin discoloration. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **Uses** This medication is used to treat acne. It may decrease the number and severity of acne pimples and promote quick healing of pimples that do develop. Tretinoin belongs to a class of medications called retinoids. It works by affecting the growth of skin cells. **How to use Tretinoin 0.025 % Topical Cream** Read the Patient Information Leaflet if available from your pharmacist before you start using this medication and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Wash your hands before applying this medication. Gently clean the affected skin with a mild or soapless cleanser and pat dry. Use your fingertips to apply a small amount of medication (about the size of a pea) in a thin layer as directed by your doctor, usually once daily at bedtime. A gauze pad or cotton swab can be used to apply the liquid. For some preparations, you should wait 20-30 minutes after cleaning your face before applying this medication. Consult the label directions, the Patient Information Leaflet, or your pharmacist if you have any questions. Use this medication on the skin only. Do not apply to the inner lip area or inside the nose/mouth. Do not apply to cut, scraped, sunburned, or eczema-affected skin. Avoid getting this medication in your eyes. If this medication gets into your eyes, flush with large amounts of water. Call your doctor if eye irritation develops. Wash your hands after using the medication to avoid accidentally getting it in your eyes. During the first few weeks of using tretinoin, your acne might appear worse because the medication is working on pimples forming inside the skin. It may take

up to 8-12 weeks to notice results from this medication. Use it regularly in order to get the most benefit from it. To help you remember, use it at the same time each day. Do not use a larger amount or use it more often than recommended. Your skin will not improve any faster, and it will increase the risk of developing redness, peeling and pain. This medication is available in different strengths and forms (such as gel, cream, solution). The best type of medication for you to use will depend on the condition of your skin and your response to therapy. Tell your doctor if your condition lasts or gets worse. Precautions Before using tretinoin, tell your doctor or pharmacist if you are allergic to it; or to vitamin A-related drugs (other retinoids such as isotretinoin); or if you have any other allergies. This product may contain inactive ingredients (such as fish proteins), which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: eczema. This medication may make you more sensitive to the sun. Limit your time in the sun. Avoid tanning booths and sunlamps. Weather extremes such as wind or cold may also be irritating to the skin. Use sunscreen daily and wear protective clothing when outdoors. Tell your doctor right away if you get sunburned or have skin blisters/redness. Wait until your skin has completely recovered from a sunburn before using tretinoin. Avoid electrolysis, waxing and chemical depilatories for hair removal on the treated areas while using this product. If you have recently used products containing sulfur, resorcinol or salicylic acid, use tretinoin with caution. Wait until the effects of such products on the skin have decreased before using tretinoin. During pregnancy, this medication should be used only when clearly needed. It may harm an unborn baby. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding. Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Cosmetics may be used, but clean skin thoroughly before applying the medication. Some cosmetics and soaps may worsen your acne. Moisturizers may be safe to use. When buying cosmetics, moisturizers or other skin care products, check the label for "non-comedogenic" or "non-acnegenic." These products are unlikely to worsen your acne. Ask your doctor or pharmacist which products are safe to use. Remember, acne is not caused by dirt. Cleaning your skin too often or too vigorously can irritate your skin and worsen acne. If you forget to use this medication, use it as soon as you remember if you remember the same day. If you don't remember until morning, wait until your next dose is scheduled. Store at room temperature. Gel preparations are flammable. Do not expose to heat or fire sources. Do not smoke during use. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Avoid using other skin products that are harsh, irritating, or drying on the treated area. These products include hair perming solutions, alcohol/lime/menthol-containing products (such as astringents, toners, shaving lotion), medicated or abrasive soaps or cleansers, soaps and cosmetics with a strong drying effect (such as alpha hydroxy acids, glycolic acid), and products containing sulfur, resorcinol, or salicylic acid. Benzoyl peroxide can be very irritating and may decrease the effectiveness of tretinoin if the two products are applied at the same time. Talk with your doctor or pharmacist about the safe use of prescription and non-prescription benzoyl peroxide products.

Side Effects

Uses

Interactions

Other Details
