

iMedix: Your Personal Health Advisor.

Typhim VI Vial

Search for medical details concerning Typhim VI intramuscular on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Redness, tenderness, hardening skin, and pain at the injection site may occur. Headache or muscle pain may also occur. If any of these effects last or get worse, tell your health care professional promptly. Rarely, some people have symptoms such as fainting, dizziness, vision changes, or ringing in the ears just after getting a vaccine injection. Tell your health care professional right away if you have any of these symptoms. Sitting or lying down may help, since these symptoms usually don't last long. Remember that this medication has been prescribed because your health care professional has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your health care professional. Contact the health care professional for medical advice about side effects. The following numbers do not provide medical advice, but in the US you may report side effects to the Vaccine Adverse Event Reporting System (VAERS) at 1-800-822-7967. In Canada, you may call the Vaccine Safety Section at Public Health Agency of Canada at 1-866-844-0018. Uses This vaccine is used to help prevent typhoid fever. Vaccines work by increasing the body's natural defense (immunity) against the bacteria that cause the infection. This vaccine is recommended for people who are traveling to areas where the infection is common (such as Africa, Asia, Central/South America), those who are in close contact to someone with an ongoing typhoid infection, and those who work with the bacteria (*Salmonella typhi*) in a lab. This vaccine is used to prevent the disease and will not treat typhoid fever or an ongoing typhoid infection. How to use Typhim VI Vial Read all vaccine information available from your health care professional before receiving the vaccine. If you have any questions, ask your health care professional. This vaccine is given by injection into the muscle of the upper arm (for adults) or into the muscle of the thigh or upper arm (for children) by a health care professional. This vaccine should be given as a single injection at least 2 weeks before possible exposure to typhoid fever. It can be given every 2 years thereafter if needed. Precautions Before receiving this vaccine, tell your health care professional if you are allergic to it; or to other vaccines; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your health care professional for more details. Before receiving this vaccination, tell your health care professional your medical history, especially of: immune system disorders (such as due to HIV infection, certain cancers such as

leukemia/lymphoma, cancer or radiation treatment), current fever/illness. During pregnancy, this vaccine should be used only when clearly needed. Discuss the risks and benefits with your health care professional. It is unknown if this vaccine passes into breast milk. Consult your health care professional before breastfeeding. Overdose Not applicable. As with any vaccine, this vaccine may not fully protect everyone who receives it. Not applicable. Store in the refrigerator. Do not freeze. Protect from light. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your health care professional. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some products that may interact with this vaccine include: drugs that weaken the immune system (including cyclosporine, tacrolimus, cancer chemotherapy, corticosteroids such as prednisone).

Side Effects

Uses

Interactions

Other Details
