## iMedix: Your Personal Health Advisor.

## **Ubrelvy**

Ubrelvy, whose generic name is ubrogepant, is an oral medication specifically designed to treat acute migraine headaches with or without aura in adults.

- ActiveIngredient:
- DosageForm:
- Dosage:
- Indications:
- Manufacturer:
- Storage:
- Market Price:
- Drug Status:

## **Description**

Side Effects Nausea or drowsiness may occur. If either of these effects lasts or gets worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada - Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. How to use Ubrelvy 100 Mg Tablet Read the Patient Information Leaflet if available from your pharmacist before you start taking ubrogepant and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth with or without food as directed by your doctor, at the first sign of a migraine. The dosage is based on your medical condition, response to treatment, and other medications you may be taking. Be sure to tell your doctor and pharmacist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). If needed, most people can take a second tablet 2 hours after the first tablet. You should not take a second tablet within 24 hours if you consume grapefruit or grapefruit juice or if you are taking certain other medications. Do not take more than 200 milligrams in a 24-hour period. Ask your doctor or pharmacist if you are not sure if or when it is safe to take a second dose. It is not known if it is safe to take this medication for more than 8 migraines within 30 days. Tell your doctor if your condition does not get better or if it gets worse. Precautions Before taking ubrogepant, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney problems, liver problems. This drug may make you drowsy. Alcohol or marijuana (cannabis) can make you more drowsy. Do not drive, use machinery, or do anything that needs alertness until you can do it safely. Limit alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is

unknown if this medication passes into breast milk. Consult your doctor before breast-feeding. Consult your pharmacist or physician. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Certain foods, beverages, or food additives (such as red wine, cheese, chocolate, monosodium glutamate) as well as lifestyle patterns such as irregular eating/sleeping habits or stress may bring on a migraine headache. Avoiding these "triggers" may help lessen migraine attacks. Consult your doctor for more details. Lab and/or medical tests (such as kidney/liver function) may be done while you are taking this medication. Keep all medical and lab appointments. Not applicable. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Other medications can affect the removal of ubrogepant from your body, which may affect how ubrogepant works. Examples include adagrasib, azole antifungals (such as ketoconazole, itraconazole), macrolide antibiotics (such as clarithromycin, erythromycin), rifamycins (such as rifampin, rifabutin), ritonavir, St.
Side Effects
Uses
Interactions

## **Other Details**