

iMedix: Your Personal Health Advisor.

Ultrasal-ER Film-Forming Solution Extended Release, With Applicator

Search for medical details concerning UltraSal-ER topical on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Slight burning, skin redness, and peeling may occur. These effects are expected. If any of these effects worsen, tell your doctor or pharmacist promptly. If your doctor has directed you to use this medication, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: spreading skin redness around the treated area, signs of infection (such as pus, milky/bloody discharge), formation of a deep sore (ulcer) at the treatment site. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. How long does it take for this medicine to take effect? Salicylic acid exerts its effects through the disruption of intercellular adhesion in the stratum corneum, leading to desquamation and enhanced removal of skin scales and plaques. The onset of action can vary depending on the concentration of salicylic acid, the specific condition being treated, and individual skin characteristics. Visible improvement in conditions such as psoriasis or wart size may be observed within 1 to 2 weeks of regular application, with more significant effects typically seen after continuous use for several weeks. How long do the effects of this medicine last? The duration of the effects of Ultrasal-ER will depend on the persistence of its application and the nature of the skin condition being treated. Continuous use maintains the keratolytic action of salicylic acid, but the effects may diminish if treatment is interrupted. For chronic conditions, ongoing treatment may be necessary to maintain improvement. Is it safe to consume alcohol while taking this medicine? There is no direct interaction between systemic alcohol consumption and the topical application of salicylic acid in Ultrasal-ER. However, alcohol can have drying effects on the skin, potentially exacerbating any dryness or irritation caused by salicylic acid. Patients should ideally maintain good skin hydration, especially when using keratolytic agents. Is this a habit forming medicine? Salicylic acid, as used in dermatological preparations like Ultrasal-ER, does not possess addictive properties and is not associated with habit formation. It is a therapeutic agent used to address specific skin conditions and does not elicit psychological

or physiological dependency. Can this medicine be taken during pregnancy? Salicylic acid is generally considered safe for topical use in limited areas and in low concentrations during pregnancy. However, high concentrations or extensive application areas should be avoided due to the theoretical risk of systemic absorption and the known effects of salicylates in pregnancy. Pregnant women should consult a healthcare provider before using Ultrasal-ER. Can this medicine be taken while breast-feeding? When used topically and in limited amounts, salicylic acid is unlikely to pose a significant risk to nursing infants. However, care should be taken to avoid application on or near the breast to prevent the infant from ingesting the product. Nursing mothers should consult a healthcare provider to determine the appropriateness of using Ultrasal-ER during breastfeeding.

Uses This medication is used on the skin to treat common skin and foot (plantar) warts. Salicylic acid helps cause the wart to gradually peel off. This medication is also used to help remove corns and calluses. This product should not be used on the face or on moles, birthmarks, warts with hair growing from them, or genital/anal warts. Salicylic acid is a keratolytic. It belongs to the same class of drugs as aspirin (salicylates). It works by increasing the amount of moisture in the skin and dissolving the substance that causes the skin cells to stick together. This makes it easier to shed the skin cells. Warts are caused by a virus. Salicylic acid does not affect the virus.

How to use Ultrasal-ER Film-Forming Solution Extended Release, With Applicator Follow all directions on the product package. If you have any questions, ask your doctor or pharmacist. This medication is for use on the skin only. To avoid irritation, do not let this medication come into contact with your eyes, nose, mouth, groin, or any broken skin. If you do get the medication in those areas, flush the area with cool water for 15 minutes. Wash hands after use. First, soak the corn, callus, or wart in warm water for about 5 minutes to soften the treatment area. Dry thoroughly. Your doctor may direct you to use a pumice stone, callus file, or emery board to carefully remove the top dead layers of skin after soaking and before applying the medication. This dead skin removal is only to help the medication work better. Do not try to rub the wart or callus off. If you are using a liquid/gel, apply a few drops or a thin coat of the medication to cover the entire wart, callus, or corn using the applicator if provided. Be careful to apply it only to the affected area and not the surrounding skin. Let dry for 5 minutes. Depending on the brand used, you may need to apply the medication twice with each treatment. Check your product package and follow the directions carefully. You may cover the area loosely with a bandage. Repeat this procedure 1 to 2 times daily for up to 2 weeks for corns and calluses and 12 weeks for warts or as directed by your doctor. If you are using a medicated pad or bandage, you may need to cut the pad so that it covers the treatment area completely but does not touch the surrounding skin. Peel off the protective covering and place the patch/bandage over the area. Leave in place according to package directions. Remove and place a new patch/bandage as directed (usually every 8 to 48 hours depending on brand). Repeat this procedure for up to 2 weeks for corns and calluses and 12 weeks for warts. Dosage is based on your medical condition, product type/brand, and response to treatment. Do not use large amounts, apply this medication more often, or use it for a longer period than directed. Your condition will not clear faster, but the chance for side effects may be increased. Use this medication regularly to get the most benefit from it. To help you remember, use it at the same time each day. If your condition lasts or gets worse, or if you think you may have a serious medical problem, get medical help right away.

Precautions Before using salicylic acid, tell your doctor or pharmacist if you are allergic to it; or to nonsteroidal anti-inflammatory drugs-NSAIDs (such as aspirin, ibuprofen, naproxen); or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: diabetes, poor blood circulation (such as peripheral vascular disease), skin infection/irritation. If you are going to have an MRI test, tell testing personnel if you are using the patch. Some patches may contain metals that can cause serious burns during an MRI. Ask your doctor whether you will need to remove your patch before the test and apply a new patch afterward, and how to do so properly. During pregnancy, this product should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Consult your doctor before breast-feeding.

Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. If your doctor has prescribed this medication for you, do not share it with others. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double

the dose to catch up. Store at room temperature. Different brands of this medication have different storage needs. Check the product package for instructions on how to store your brand, or ask your pharmacist. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor’s approval.

Side Effects

Uses

Interactions

Other Details
