

iMedix: Your Personal Health Advisor.

Ustekinumab Solution

Search for medical details concerning ustekinumab intravenous on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Bruising, itching, pain, redness, swelling, or hardening of the skin at the injection site may occur. Injection site reactions usually go away after 1 or 2 days. Headache, back pain, or sinus/throat pain may also occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: cough that doesn't go away, shortness of breath. This medication can affect your immune system. It can lower your body's ability to fight an infection. You may be more likely to get serious infections, such as lung infections, bone/joint infections, skin infections, sinus infections, or bowel/gallbladder infections. It may also be harder to fight an infection you already have. Tell your doctor right away if you develop any signs of an infection, such as nausea/vomiting that doesn't stop, worsening redness/swelling/tenderness at the injection site after 2 days, fever/chills, cold/flu symptoms, painful/frequent urination, unusual vaginal discharge/burning/itching/odor, or severe stomach pain. () Ustekinumab may cause a rare (sometimes fatal) condition called PRES (posterior reversible encephalopathy syndrome). Get medical help right away if you develop headache that doesn't go away, seizures, sudden vision changes, mental/mood changes (such as confusion). There is a rare risk of developing cancer (including skin cancer) due to this medication. Discuss the risks and benefits of treatment with your doctor. Tell your doctor right away if you develop symptoms such as unusual lumps/growths, unusual skin changes (including a sore that does not heal or a change in the size/shape/color of a mole), swollen glands, unexplained weight loss. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. **How to use Ustekinumab Solution** Read the Medication Guide and the Instructions for Use provided by your pharmacist before you start using ustekinumab and each time you get a refill. If you have any questions, consult your doctor or pharmacist. Before starting this drug, you should take a tuberculosis (TB) skin test to check for a type of tuberculosis that may not be causing any symptoms (latent TB). If you are diagnosed with TB, to prevent a serious TB infection you must first be

treated for it before you start ustekinumab. For the treatment of psoriasis or psoriatic arthritis, this medication is given by injection under your skin as directed by your doctor. You will receive one dose, followed by a second dose 4 weeks later. Then this medication is given every 3 months. For the treatment of Crohn's disease or ulcerative colitis, the first dose of this medication is given by injection into a vein by a health care professional. The medication is then given by injection under your skin every 8 weeks as directed by your doctor. The dosage is based on your weight, medical condition, and response to treatment. If you are giving this medication to yourself at home, learn all preparation and usage instructions from your health care professional. Do not shake the solution. The solution is normally colorless to light yellow. It may contain a few small white particles of protein. Before using, check this product visually for other particles or discoloration. If either is present, do not use the liquid. Learn how to store and discard medical supplies safely. Do not reuse syringes. When given under the skin, this medication should be injected in either the upper arms, buttock, thighs, or abdomen. It is important to change the location of the injection site with each dose to avoid problem areas under the skin. Choose a different injection site with each dose. Do not inject into skin that is irritated, sore, or infected. Use this medication regularly to get the most benefit from it. It may help to mark your calendar with a reminder. Tell your doctor if your condition does not improve or if it worsens. Precautions Before using ustekinumab, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients (such as natural rubber/latex found in the needle cover on the prefilled syringe), which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: current/recent/returning infection (including hepatitis and tuberculosis), cancer. Ustekinumab can make you more likely to get infections or may make current infections worse. Do not have immunizations/vaccinations without the consent of your doctor, and avoid contact with people who have recently received oral polio vaccine or flu vaccine inhaled through the nose. Avoid receiving BCG vaccines for one year after completing treatment with ustekinumab. Also, stay away from anyone who has an infection that may easily spread (such as chickenpox, COVID-19, measles, flu). Talk to your doctor if you have been exposed to an infection or for more details. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. This drug may pass into breast milk. Consult your doctor before breast-feeding. Consult your pharmacist or physician. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. Lab and/or medical tests (such as skin exams) should be done while you are using this medication. Keep all medical and lab appointments. Consult your doctor for more details. It is important to get each dose of this medication as scheduled. If you miss a dose, ask your doctor or pharmacist right away for a new dosing schedule. Do not double the dose to catch up. Different brands of this medication have different storage needs. Check the product package for instructions on how to store your brand, or ask your pharmacist. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Side Effects

Uses

Interactions

Other Details
