

# iMedix: Your Personal Health Advisor.

## Veltassa 8.4 Gram Oral Powder Packet

---

Search for medical details concerning Veltassa oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
  - **DosageForm:**
  - **Dosage:**
  - **Indications:**
  - **Manufacturer:**
  - **Storage:**
  - **Market Price:**
  - **Drug Status:**
- 

### Description

Side Effects Constipation or diarrhea may occur. If either of these effects lasts or gets worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Patiromer may cause a low magnesium blood level. Your doctor will check your level and prescribe a magnesium supplement if your level is low. Tell your doctor right away if you have: symptoms of a low magnesium blood level (such as muscle cramps, irregular heartbeat, mental/mood changes like confusion). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. How to use Veltassa 8.4 Gram Oral Powder Packet Take this medication by mouth with or without food as directed by your doctor, usually once daily. The dosage is based on your medical condition, potassium lab test results, and response to treatment. Do not heat patiromer or add it to hot foods or liquids. Mix this medication right before taking it. Open the packet and mix the powder as directed with liquid (such as water, apple juice) or soft food (such as applesauce, pudding, yogurt). If using liquid to mix, the powder does not dissolve so the liquid mixture will look cloudy. If any powder is left in the glass after drinking the mixture, add more liquid, stir, and drink it right away to make sure you take the whole dose. If using food to mix, make sure to eat all of the mixture to make sure you take the whole dose. Patiromer may decrease your absorption of other medications, which may make them work less well. Ask your pharmacist about the other medications you take. You may need to take patiromer at least 3 hours before or 3 hours after certain medications. Precautions Before taking patiromer, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: problems with your bowel movement (such as severe constipation, intestinal/bowel blockage, dry hard stool/impaction, slowed bowel after stomach/intestinal surgery), low magnesium blood level. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and

herbal products). Older adults may be more sensitive to the side effects of this drug, especially constipation or diarrhea. During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. This medication does not pass into breast milk. Consult your doctor before breast-feeding. Consult your pharmacist or physician. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: muscle cramps, weakness, irregular heartbeat. Do not share this medication with others. Lab tests (such as magnesium and potassium blood levels) must be done while you are taking this medication. Keep all medical and lab appointments. If your doctor has prescribed a special diet for you, follow it. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store in the refrigerator. If stored at room temperature, use it within 3 months of taking it out of the refrigerator. Avoid high heat. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

---

---

## **Side Effects**

---

---

## **Uses**

---

---

---

---

---

---

## **Interactions**

---

---

## **Other Details**

---

