

# iMedix: Your Personal Health Advisor.

## Viagra Soft (Sildenafil)

---

Viagra Soft offers sildenafil citrate in a pliable tablet formulation, potentially allowing for quicker absorption or easier administration for men managing erectile difficulties, especially those who prefer an alternative to conventional hard pills. Its design often aims to enhance user convenience and may offer a different onset experience.

- **ActiveIngredient:** Sildenafil Citrate
  - **DosageForm:** Soft tablet
  - **Dosage:** 50mg and 100mg
  - **Indications:** Erectile dysfunction
  - **Manufacturer:** Pfizer Inc.
  - **Storage:** Keep at controlled room temperature, ideally between 15°C and 30°C (59°F to 86°F), shielded from excessive moisture, heat, and direct light.
  - **Market Price:** 0.91
  - **Drug Status:** Prescription Only
- 

## Description

General information Erectile dysfunction is a pathology which characterising premature ejaculation, as well as a short erection or its absence. This disease is the result of many reasons, for example, the negative influence of hormonal drugs, bad habits, malnutrition, sedentary lifestyle, stress, fatigue, lack of sleep and others. Hereby we present you a popular drug for potency – Viagra Soft, which aims at sexual life improvement. Safety and reliability of this drug have big popularity and recognition among many men. Action of the drug First of all, it is worth noting its availability, more rapid onset of action which you can compare with the classical type of drug. This action is contributes by the specific formula of this pill, namely, the possibility to dissolve it, which results in a faster flow of the active substance into the blood. Also it ought to notice the gentler, delicate impact of the medication on the body. It is due to the absence of an irritant factor on the gastrointestinal tract. Furthermore this product has a pleasant taste. Viagra Soft is an excellent product for potency stimulating. This remedy is for sale in the form of soft drugs, which, as a rule, has a pleasant taste. The main difference of this medication comparing to the classic remedy is the possibility to dissolve in mouth, leading to the assimilation process reduction to 20 minutes. How long does it take for this medicine to take effect? The onset of action of Viagra Soft, containing sildenafil citrate, typically occurs within 15 to 30 minutes after administration. Sildenafil is a phosphodiesterase type 5 (PDE5) inhibitor that enhances the effects of nitric oxide, leading to relaxation of smooth muscles in the corpus cavernosum of the penis and increased blood flow, resulting in an erection when sexually stimulated. How long do the effects of this medicine last? The effects of Viagra Soft usually last for approximately 4 to 6 hours after ingestion. However, individual response may vary depending on factors such as dosage, metabolism, and presence of underlying medical conditions. Is it safe to consume alcohol while taking this medicine? While moderate alcohol consumption is unlikely to significantly affect the pharmacokinetics of sildenafil, excessive alcohol intake may increase the risk of adverse effects such as dizziness, hypotension, and impaired judgment. Patients should exercise caution when consuming alcohol while taking Viagra Soft and avoid excessive drinking to minimize the potential for adverse reactions. Is this a habit forming medicine? Viagra Soft is not considered a habit-forming medicine. It does not contain substances with addictive properties, and discontinuation of treatment typically does not result in withdrawal symptoms. However, it is essential to use

Viagra Soft only as prescribed by a healthcare professional to ensure its safe and effective use. Can this medicine be taken during pregnancy? Viagra Soft is not indicated for use in women, including during pregnancy. It is specifically prescribed for the treatment of erectile dysfunction in men and should not be used by pregnant individuals. Women who are pregnant or planning to become pregnant should not handle Viagra Soft tablets due to the potential risk of adverse effects on the fetus. Can this medicine be taken while breast-feeding? Due to the lack of adequate safety data, Viagra Soft should not be used while breastfeeding. Sildenafil may pass into breast milk and pose potential risks to the nursing infant. Women who are breastfeeding should discuss alternative treatment options with their healthcare provider to ensure the safety of their child while receiving appropriate treatment for erectile dysfunction.

**Advantages** Also, the drug practically has no side effects and it is effective at solving problems with erectile dysfunction. Unlike other drugs which exist in the market for potency stimulation, this drug you can find in the form of soft pills, which you can both dissolve in mouth or intake like regular tablets. Due to such a property, the speed of impact on the body is significantly reducing from 40-50 minutes to 15-20 minutes. The main ingredient of this drug is Sildenafil, which does not affect the hormones, but reduces the speed of phosphodiesterase enzymes action. You should remember that sexual stimulation results in an erection. Sildenafil increases blood flow to the penis, which results and provides a lasting effect that lasts up to 4-5 hours. This drug is made successfully helps men who have problems with sexual function, regardless of age and causes of its occurrence. Moreover, the drug has a powerful antibacterial and antiviral effect and protects the reproductive system against inflammation and the development of infectious diseases.

**Precautions** You should take this drug with caution by persons with individual intolerance to any substances which also Viagra includes. Also, special attention you should pay if you have problems with the cardiovascular, liver and kidneys systems. In addition, men over the age of 65 years before taking the product have to undergo a medical examination to avoid the possible complications occurrence. This medication is not for use for people younger 18 y.o. and women. We should note that this medication has contraindications in case of increased sexual activity due to health reasons. Also you can't take this drug if you had a heart attack recently or stroke. You should not use it in cardiac, renal or hepatic failure, retina hereditary pathology, also with penile injuries and chronic gastrointestinal diseases.

**Dosage** The maximum daily dosage of the remedy is 100 mg. For most men, this remedy has a good effect in a dose of 50 mg. With individual intolerance to separate components of the drug, you can reduce the dose to 25 mg, or in case of a weak susceptibility – increase the dosage to 100 mg. You should get a doctors examination for the final decision on the dosage. He will take into account the state of your health, as well as possible complications after illness. But there are general rules for taking this drug: you should take it about half an hour prior sexual intercourse; put the soft drug under the tongue and dissolve or chew. The erection onset is possible only with natural arousal. Without erotic stimulation, the medication cannot cause any spontaneous erection. The effect of the drug begins approximately in 30 minutes after administration. Of course, the particular characteristics of the body of each individual man are important here. Remember that after taking a large amount of food, the effect of Viagra Soft may begin later.

**How to take?** For the sooner effect of the product action specialists suggest to dissolve or chew the soft drug 15-30 minutes before sexual act. The duration of the drug action is about 4-5 hours. The optimal dosage for a single dose is 50 mg or 1 soft drug. Doctors recommend to take drugs before meals. In this case, the impact of the product comes faster. The fatty foods prevent the drug's high-quality absorption in the the blood stream and slows down the impact of this product. In this case it is better to prefer cereal dishes, fruits and vegetables. At the same time this drug is not good to mix with the medicines, which include nitric oxide donators, alpha-blockers, nitrates as well as inhibitors of other species.

**Side-effects** Viagra Soft does not give an addiction to it and never affects the sperm. Side effects occur very rarely. In most cases, they can occur if there was an overdose of the product. The patient may complain of nausea, nasal congestion or headache. During clinical investigations, depending on individual doses selection, most common were next side reactions: heart and vessels system – flushing, cephalgia, dizziness; organs of the digestive system – indigestion; respiratory system – nasal blockage; sensory organs – altered vision (impaired visual clarity and others).

**Overdose** According to investigation which has been performed with participation of healthy volunteers using a single dose in the amount of 200 mg, there were manifested some adverse reactions which were similar to those occurred when Viagra Soft was used in smaller amounts. Standard symptomatic treatment should be taken in the case of overdose followed by medical help.

**Interactions** Viagra Soft should not be combined with drugs that include all types of nitrates, as well as other inhibitors types, and nitrogen donators. Such medications

can increase the Sildenafil concentration in the blood plasma. It should be remembered that the substance co-administered with drugs containing these components is hazardous to health. When treating chronic diseases with other groups of medical agents, it is recommended to consult with your doctor. Alcohol interaction Viagra Soft should never be used simultaneously with alcoholic beverages. Drug and alcohol combination can cause unforeseen phenomena, greatly enhance the adverse reactions manifestation; provoke intoxication and a sharp deterioration in the person's well-being.

---

---

**Side Effects**

---

**Uses**

---

---

---

---

---

---

**Interactions**

---

- - 
  - 
  -
- 

**Other Details**

---

---