

# iMedix: Your Personal Health Advisor.

## Vidalista 80mg Yellow

---

Vidalista 80mg Yellow delivers a substantial dose of tadalafil intended for men requiring potent intervention for erectile dysfunction, identifiable by its distinct yellow color to aid recognition and safe usage. This formulation provides the extended duration effect associated with tadalafil in a high-strength format.

- **ActiveIngredient:** Tadalafil
  - **DosageForm:** Oral tablet
  - **Dosage:** 80 mg
  - **Indications:** Utilized in treating male erectile dysfunction (ED)
  - **Manufacturer:** Varies (Centurion Laboratories commonly associated; confirm via packaging)
  - **Storage:** Keep below 30°C (86°F), away from moisture and direct sunlight. Maintain in original container. Store out of children's view and access.
  - **Market Price:** 0.75
  - **Drug Status:** Prescription Only
- 

## Description

What is Vidalista 80mg Yellow? Vidalista 80mg Yellow is a pharmaceutical preparation indicated for adult men experiencing difficulty with erectile function (erectile dysfunction). It contains a significant 80 mg strength of Tadalafil, its sole active ingredient. This specific variant is identifiable by its distinctive yellow coloration, which helps differentiate it from other strengths or medications. Mechanism of Action The therapeutic activity of tadalafil is achieved through its inhibition of the phosphodiesterase type 5 (PDE5) enzyme. By impeding PDE5, tadalafil helps preserve levels of cyclic guanosine monophosphate (cGMP) within the penile smooth muscle tissues when sexual stimulation occurs. This biochemical process facilitates the relaxation of blood vessels supplying the penis, promoting the increased blood flow essential for achieving and maintaining an erection. The substantial 80 mg dose is intended to provide a pronounced effect within this mechanism for certain individuals. The potential for erectile assistance can last for an extended period, often cited as up to 36 hours. Brand vs. Generic Status Marketed under the name Vidalista 80mg Yellow, this product is a branded generic formulation. It employs tadalafil, the same active pharmaceutical agent found in the reference drug, Cialis®. However, it is presented under its own trade name and with its characteristic yellow appearance, differentiating it from the originator product and other generic versions. Available Forms This medication is supplied as oral tablets, each containing a fixed dose of 80 milligrams of tadalafil. These tablets are specifically colored yellow and are intended for oral consumption according to a healthcare provider's instructions. Unique Features Substantial Dosage: Features a high 80 mg dose of tadalafil, aimed at individuals who may require a more potent treatment effect under medical supervision. Visual Identification Aid: The distinct yellow color serves as a visual cue, potentially aiding in patient identification, adherence, and preventing confusion with lower doses or different medications. Extended Efficacy Window: Consistent with tadalafil, it offers a prolonged duration (up to 36 hours) during which achieving an erection upon sexual arousal may be facilitated. Targeted High-Strength Option: Represents a specific high-dose choice within the Vidalista line, intended for particular patient needs as assessed by a doctor.

---

## General Instructions

**How to Take Vidalista 80mg Yellow Administration:** Consume one yellow Vidalista 80mg tablet by mouth using a glass of water. It is important to swallow the tablet without altering it – do not break, crush, or chew the medication. **Food Considerations:** This medication may be ingested with or without food. However, be aware that consuming a meal particularly high in fat content close to the time of administration might slightly prolong the time until the drug takes effect. **Timing Guidelines Optimal Timing:** Aim to take the tablet approximately 30 to 60 minutes before you intend to be sexually active. Remember that achieving an erection requires concurrent sexual stimulation. **Duration of Effect:** The potential for Vidalista 80mg Yellow to assist with erectile function can span a significant period, potentially extending up to 36 hours after taking the tablet. **Frequency Maximum Dosage:** Due to the substantial 80 mg strength, it is absolutely essential to adhere to a maximum limit: do not ingest more than a single Vidalista 80mg Yellow tablet within any 24-hour interval. **Missed Dose Situational Use:** Vidalista 80mg Yellow is designed for use on an ‘as-needed’ basis preceding sexual activity, not on a fixed daily schedule. Therefore, if you forget to take it at a planned moment, simply administer the dose when needed, ensuring the 24-hour minimum gap since the last intake is maintained. **Storage Guidelines Conditions:** Keep your Vidalista 80mg Yellow tablets stored at a controlled room temperature, generally advised as below 30°C (86°F). **Protection:** Retain the tablets within their original blister pack until ready for consumption to shield them from detrimental effects of moisture and light. Critically, ensure the medication is stored in a location inaccessible to children and pets.

---

## Side Effects

Utilizing Vidalista 80mg Yellow might lead to adverse effects in certain users, as can occur with any medication. The significant 80 mg tadalafil dose could potentially influence the frequency or intensity of these reactions compared to lower strengths. **Common Side Effects:** These are encountered relatively often but are generally not severe and may subside as your system adapts. **Head Discomfort:** Experiencing a headache is fairly typical. Ensuring adequate fluid intake might offer some relief. **Facial Warmth/Redness:** A temporary feeling akin to blushing or warmth spreading across the face. **Nasal Blockage:** A sensation of having a stuffy or obstructed nose. **Digestive Unease:** Mild symptoms like indigestion or a slightly upset stomach following tablet intake. **Muscle or Back Discomfort:** Aches sometimes arise, often developing 12-24 hours after taking the dose and usually fading within 48 hours. Your doctor might suggest simple pain relief if necessary. **Less Common Side Effects:** These effects are noted less frequently but are still possibilities. **Dizziness Sensation:** Feeling lightheaded or experiencing a loss of balance. Avoid tasks needing full concentration if affected. **Visual Adjustments:** Transient issues such as slightly unfocused vision or subtle changes in color perception. **Increased Heartbeat:** Awareness of your heart beating faster than its usual pace. **Eye Discomfort:** Feelings of pain or visible redness within the eyes. **Serious Side Effects:** These situations are rare but necessitate immediate medical attention. Stop using Vidalista 80mg Yellow and get emergency help right away if you observe: **Priapism (Extended Erection):** An erection persisting painfully for more than four consecutive hours. This demands urgent treatment to prevent potential irreversible penile injury. **Sudden Sight Reduction:** An abrupt decrease or total loss of vision impacting one eye or both (could signal NAION). **Abrupt Hearing Deficit:** A sudden drop in hearing ability, potentially associated with ringing noises (tinnitus) or feelings of dizziness. **Cardiac Warning Signs During Intimacy:** Experiencing chest pain, acute dizziness, or persistent nausea while engaging in sexual activity may indicate a serious heart issue, particularly if pre-existing cardiac conditions are present. **Major Hypersensitivity Event:** Symptoms might include a significant rash, hives, severe itching, respiratory distress, or notable swelling involving the face, mouth, tongue, or throat. **Management Note:** Should any side effect feel intense, fail to resolve, or cause significant worry, consulting your healthcare professional for tailored advice is crucial. Do not attempt to change your medication regimen without medical input.

---

## Uses

**Approved Medical Application** Vidalista 80mg Yellow is employed in the therapeutic management of erectile dysfunction (ED) in the adult male population. This condition is understood as the recurrent

inability to generate or uphold a penile erection suitable for satisfactory sexual relations. **How It Addresses This Condition** **Potent Vascular Enhancement:** The substantial 80mg tadalafil load in Vidalista 80mg Yellow functions by powerfully obstructing the phosphodiesterase type 5 (PDE5) enzyme within penile tissues. This intervention amplifies the natural signaling pathway involving cGMP that occurs during sexual arousal, resulting in pronounced relaxation of arterial smooth muscles and facilitating the robust inflow of blood needed for a firm erection. **Extended Opportunity for Intimacy:** A hallmark benefit derived from tadalafil, including this high-strength yellow variant, is the significantly prolonged timeframe (potentially reaching 36 hours) during which erectile response to sexual stimuli may be enhanced. **High-Intensity Effect:** The 80 mg dosage is specifically intended to elicit a strong physiological response, potentially benefiting individuals for whom standard doses of tadalafil have proven insufficient, following careful medical evaluation. **Distinct High-Dose Option:** This formulation represents a clearly identifiable high-strength choice within the product line for managing ED. **Important Considerations** **Focus on Symptom Management:** Users should understand that Vidalista 80mg Yellow alleviates the physical symptoms of ED but does not resolve or cure underlying contributors, which can range from vascular disease and neurological issues to psychological factors or medication side effects. **Sexual Stimulation is Imperative:** The medication aids the body's natural response mechanism; it does not create an erection in the absence of genuine sexual excitement or arousal. **High Dosage Requires Justification:** It's vital to recognize that the 80 mg strength represents a significant therapeutic dose, necessitating careful physician assessment regarding its appropriateness and safety before initiation; it is generally not a first-line treatment choice. **Value of Holistic Health:** Optimal results in managing ED are often achieved when pharmacotherapy like Vidalista 80mg Yellow is integrated with broader health strategies, such as addressing cardiovascular risk factors, adopting a healthy lifestyle, and managing related conditions like diabetes or hypertension. **Color as an Identifier:** The distinct yellow appearance of this specific 80mg formulation can serve as a helpful visual cue for patients and caregivers, assisting in differentiating it from lower strengths or other medications, thereby contributing to safer medication handling and adherence practices.

---

## **Safety advice**

### **Interactions Alcohol:**

- Use with caution
- Combining Vidalista 80mg Yellow with alcohol can markedly amplify the potential for adverse reactions like dizziness or sudden blood pressure drops due to the high tadalafil dose. Exercising significant restraint regarding alcohol consumption is therefore strongly recommended when using this medication.

### **Interactions Other Medications:**

- Consult your doctor
- The substantial 80 mg tadalafil content in Vidalista 80mg Yellow creates a heightened potential for clinically significant interactions with numerous drugs, especially nitrates (absolutely forbidden), alpha-blockers, potent enzyme modulators, and certain antihypertensives. Providing your clinician with an exhaustive list of all concurrent medications and supplements is critical for safety.

### **Special Groups Pregnancy:**

- Unsafe
- This medication is exclusively intended for managing male erectile dysfunction and must never be administered to women, with absolute contraindication during pregnancy.

### **Special Groups Breastfeeding:**

- Unsafe
- Vidalista 80mg Yellow is not authorized for female use under any circumstances, including during the period of lactation.

### **Special Groups Elderly:**

- Use with caution
- Individuals aged 65 and over may exhibit heightened sensitivity to tadalafil's effects, particularly at the substantial 80 mg dose, necessitating careful evaluation by a physician regarding suitability and potential risks like exaggerated blood pressure drops.

### **Special Groups Children:**

- Unsafe
- Administration of Vidalista 80mg Yellow to any individual below the age of 18 is strictly contraindicated, as its safety and effectiveness in minors have not been investigated.

### **Effects on Activities Driving:**

- Use with caution
- As tadalafil can occasionally induce dizziness or visual disturbances, users should exercise caution and understand their personal response before operating a motor vehicle after taking Vidalista 80mg Yellow.

### **Effects on Activities Operating Machinery:**

- Use with caution
- Until familiar with how Vidalista 80mg Yellow affects them personally, individuals should be cautious when engaging in tasks demanding full alertness and coordination, such as operating potentially hazardous machinery.

---

## **Concerns**

**Key Concerns When Considering Vidalista 80mg Yellow Significant Hemodynamic Impact:** The potent 80 mg tadalafil concentration in Vidalista 80mg Yellow can induce substantial changes in blood pressure. This warrants extreme caution, particularly in individuals with underlying cardiovascular vulnerabilities, those using other blood pressure medications, or anyone susceptible to dizziness or fainting spells.

**Theoretical Priapism Risk:** While priapism (a prolonged, painful erection over 4 hours) is an inherently rare side effect of tadalafil, the use of a very high 80 mg dose theoretically might slightly elevate this risk profile. Awareness and readiness to seek immediate medical help are crucial.

**Potential for Sensory Adverse Events:** Although uncommon, sudden alterations in vision (like NAION) or hearing have been linked to PDE5 inhibitors. The high systemic exposure from an 80 mg dose underscores the importance of promptly reporting any such sudden sensory changes to a healthcare provider.

**Metabolic Load on Organs:** Processing a substantial 80 mg dose of tadalafil places a significant metabolic demand on the liver and kidneys. Individuals with pre-existing moderate to severe impairment of these organs face a higher risk of drug accumulation and intensified side effects, potentially making this dose unsuitable.

**Color-Coding and Adherence:** While the yellow color aids identification, the high potency necessitates meticulous adherence to the prescribed regimen. Accidental double-dosing or confusion with lower strengths could lead to serious adverse effects.

**When Not to Use Vidalista 80mg Yellow Absolute Contraindications:** Use is unequivocally forbidden for individuals currently taking any form of organic nitrate medication (due to catastrophic hypotension risk). Other fundamental contraindications include a history of severe hypersensitivity to tadalafil, diagnosis of specific unstable cardiovascular conditions (recent stroke/heart attack, severe heart failure, uncontrolled arrhythmias), previous experience with NAION-related vision loss, and diagnosis of severely compromised liver function. Use in end-stage renal disease is also generally precluded.

**Critical Reminders High Dose Requires Expert Oversight:** Stress that Vidalista 80mg Yellow represents a maximum strength formulation, not intended as initial ED therapy. Its use demands careful evaluation and ongoing supervision by a qualified healthcare professional.

**Yellow Color for Safety:** Utilize the distinctive yellow color as a key identifier to prevent medication errors, especially distinguishing it from lower tadalafil doses or other drugs. Ensure correct medication selection every time.

**Elevated Counterfeit Risk:** Be particularly vigilant about the source of high-strength generic medications like Vidalista 80mg Yellow. Counterfeit versions are prevalent and may contain dangerous inconsistencies or contaminants. Obtain only via prescription from reputable suppliers.

---

## Warnings

Critical Health Authority Precautions Regulatory bodies issue significant cautions regarding tadalafil, the active substance in Vidalista 80mg Yellow, particularly relevant given its high dose: Nitrate Co-use Danger: An overriding warning concerns the potential for a precipitous, potentially fatal decline in systemic blood pressure should tadalafil be ingested concurrently with any organic nitrate preparation. Mandatory Cardiac Evaluation: Due to tadalafil's hemodynamic effects, a thorough assessment of a patient's cardiovascular fitness is considered essential prior to initiating therapy, especially at the 80 mg strength. Rare Vision Loss Phenomenon (NAION): Although infrequent, cases of non-arteritic anterior ischemic optic neuropathy (NAION), resulting in sudden vision compromise, have been documented following the use of PDE5 inhibitors like tadalafil. Absolute Contraindications Usage of Vidalista 80mg Yellow is strictly prohibited under the following specific conditions: Simultaneous Nitrate Administration: Taking this medication alongside any form of nitrate drug (e.g., nitroglycerin, isosorbide compounds) or recreational nitrites ("poppers") is absolutely forbidden due to the risk of profound hypotension. Documented Tadalafil Hypersensitivity: Individuals known to have experienced a significant allergic reaction to tadalafil or any non-active component of the Vidalista 80mg Yellow tablet must avoid its use. Defined Severe Cardiovascular States: Patients with markedly unstable cardiac conditions – such as very recent (within 6 months) myocardial infarction or stroke, certain classifications of advanced heart failure, uncontrolled severe arrhythmias, or resting blood pressure significantly outside normal ranges (severe hypo- or hypertension) – are contraindicated. Urgent Precautions The development of any of the following scenarios after consuming Vidalista 80mg Yellow necessitates immediate cessation of the drug and urgent medical consultation: Priapism Episode: An erection persisting for a duration exceeding four hours (painful or otherwise). This is a medical emergency requiring prompt treatment to avoid potentially irreversible damage to penile structures. Abrupt Sensory Impairment: Any sudden decrease or complete loss of vision (in one or both eyes) or hearing (possibly with associated tinnitus or dizziness). Discontinue use immediately and seek emergency medical evaluation. Potential Cardiac Event Symptoms: Experiencing symptoms like chest discomfort, acute dizziness or fainting, or prolonged nausea during or immediately following sexual activity. These could indicate a serious cardiovascular reaction needing urgent medical review.

---

## Dosage

Standard Dosing Guidelines Prescribed Amount: Vidalista 80mg Yellow delivers a substantial 80 milligrams of tadalafil in each tablet. The typical instruction involves taking a single 80 mg tablet orally, when required, before engaging in sexual activity. It must be understood that this significant dosage strength is generally reserved for specific clinical situations determined by a healthcare provider and is not a routine starting point for ED management. Maximum Frequency: Owing to the high drug load and tadalafil's extended duration of action (up to 36 hours), a strict limit applies: do not consume more than one Vidalista 80mg Yellow tablet within any single 24-hour cycle. Dose Adjustments Renal Impairment: Profound caution is mandated. For individuals with mild to moderate kidney difficulties, utilizing such a high dose may necessitate close observation or might be inappropriate. Vidalista 80mg Yellow is typically strongly discouraged or contraindicated for patients suffering from severe kidney disease or those undergoing dialysis, as tadalafil elimination is impaired, potentially leading to dangerously elevated drug concentrations and heightened side effect risk. Physician guidance is paramount. Hepatic Impairment: Similar vigilance applies regarding liver function. Persons with mild to moderate liver problems should only consider this high dose under rigorous medical oversight, if at all. Vidalista 80mg Yellow is generally contraindicated (must not be used) by individuals diagnosed with severe hepatic failure, given the liver's crucial role in metabolizing tadalafil. Elderly Patients: Senior individuals (customarily defined as 65 years or older) might react more intensely to tadalafil's effects and side effects. The 80 mg dose carries potentially greater risks (like orthostatic hypotension) within this demographic. Dosage suitability must be carefully determined by a medical professional, weighing overall health status and tolerance. Missed Dose Since Vidalista 80mg Yellow is formulated for administration on an 'as-needed' basis before sexual relations, a regular dosing timetable isn't

followed, making the concept of a ‘missed dose’ largely inapplicable. Simply administer a dose when the need arises, while strictly observing the 24-hour interval rule since any prior intake. Overdose Management Symptoms: Consuming more than the directed dose of Vidalista 80mg Yellow constitutes an overdose. Potential manifestations typically represent magnified versions of known side effects, possibly including intense headache, dangerously prolonged and painful erection (priapism), severe dizziness or fainting (resulting from significant hypotension), marked muscle soreness, pronounced flushing, or considerable nausea/vomiting. Action: Should an overdose be suspected, securing immediate emergency medical aid is non-negotiable. Contact emergency response services or a poison control center promptly. Management will focus on providing supportive care addressing the specific symptoms presented (e.g., interventions for priapism, blood pressure support).

---

## Interactions

**Drug-Drug Interactions Nitrates (Absolute Preclusion):** The concurrent use of Vidalista 80mg Yellow with any form of organic nitrate medication (typically for angina, e.g., nitroglycerin, isosorbide mononitrate/dinitrate) or recreational nitrite inhalants (“poppers”) constitutes a non-negotiable prohibition. This combination can trigger a rapid, severe, and potentially lethal drop in blood pressure. **Alpha-Blockers and Other Antihypertensives:** Caution is strongly advised when combining this high-dose tadalafil formulation with alpha-adrenergic blockers (used for BPH/hypertension, e.g., doxazosin) or other agents intended to lower blood pressure. Both drug types possess hypotensive effects, and their simultaneous use creates a compounded risk of symptomatic low blood pressure (dizziness, fainting). Physician oversight regarding dosing or timing may be essential. **Potent Inhibitors of CYP3A4 Metabolism:** Medications known to strongly inhibit the CYP3A4 enzyme pathway (such as ketoconazole, itraconazole, clarithromycin, HIV protease inhibitors like ritonavir) can substantially impede the breakdown of tadalafil. This leads to markedly increased bioavailability and heightened potential for dose-related side effects from the 80 mg strength. Use alongside such inhibitors requires extreme caution or may be contraindicated per medical advice. **Potent Inducers of CYP3A4 Metabolism:** Conversely, potent CYP3A4 inducers (e.g., rifampin, carbamazepine, phenytoin, phenobarbital) can significantly accelerate the metabolic clearance of tadalafil. This action can result in reduced systemic exposure, potentially subverting the intended therapeutic effect of Vidalista 80mg Yellow. **Other PDE5 Inhibitors or Riociguat:** The simultaneous administration of Vidalista 80mg Yellow with other phosphodiesterase type 5 inhibitors (e.g., sildenafil, vardenafil) or the guanylate cyclase stimulator riociguat is strictly discouraged. Such combinations offer no proven benefit and substantially amplify the risk of excessive blood pressure reduction and other overlapping adverse effects. **Drug-Food Interactions Meals High in Fat Content:** While tadalafil absorption is generally less sensitive to food compared to some alternatives, consuming a meal particularly rich in fat immediately before taking Vidalista 80mg Yellow might marginally delay the absorption rate and the time required to achieve peak effect for some individuals. **Grapefruit and Related Citrus:** Grapefruit juice (and potentially grapefruit itself) contains compounds that can disrupt the CYP3A4 metabolic pathway in the gut wall, potentially resulting in elevated plasma concentrations of tadalafil. Substantial consumption should likely be avoided to prevent an increased risk profile. **Alcoholic Beverages:** Exercise significant prudence regarding alcohol intake. Combining substantial alcohol amounts with the potent 80 mg tadalafil dose can exacerbate susceptibility to side effects like headache, dizziness, increased heart rate, and orthostatic hypotension (postural dizziness/fainting). Alcohol can also independently impair erectile capacity. **Drug-Supplement/Herb Interactions St. John’s Wort (Hypericum perforatum):** This botanical supplement acts as a CYP3A4 inducer and could potentially decrease the overall effectiveness of tadalafil by speeding its elimination. Discuss its use with your doctor. **Supplements with Hypotensive Activity:** Exercise caution with supplements known or suspected to lower blood pressure (e.g., high-dose fish oil, certain amino acids like L-arginine), as they might theoretically have additive effects with Vidalista 80mg Yellow. **Impact of Interactions** The potential ramifications of interactions involving Vidalista 80mg Yellow span a spectrum, from potentially neutralizing the medication’s benefit (due to induced metabolism) to significantly magnifying the risk of serious adverse outcomes (like profound hypotension with nitrates or amplified side effects with CYP3A4 inhibitors). Some interactions necessitate urgent medical evaluation. **Critical Reminders Comprehensive Disclosure is Paramount:** Given

the potency of Vidalista 80mg Yellow, comprehensive transparency with your physician regarding all ingested substances—prescriptions, over-the-counter products, vitamins, herbs, recreational drugs—is paramount to preempt dangerous interactions. Magnified Counterfeit Interaction Risk: The risk of unpredictable, harmful interactions is particularly magnified with counterfeit high-strength medications like Vidalista 80mg Yellow, which may harbor undisclosed or incorrect ingredients reacting dangerously with other substances.

---

- - 
  - 
  -
- 

## Other Details

**Appearance** Vidalista 80mg Yellow tablets are distinguished by their characteristic yellow color. They are typically oral, film-coated tablets, but the precise shade of yellow, shape (e.g., round, almond), and any specific markings or engravings can vary depending on the manufacturing batch and source (refer to packaging; Centurion Laboratories is a common manufacturer). **Storage After Opening** To maintain optimal integrity, each Vidalista 80mg Yellow tablet should remain sealed within its original blister strip until immediately prior to administration. This practice protects the high-dose tablet from degradation due to environmental factors like humidity and light. Storing them loose or in multi-day pill containers is not advised. **Disposal** Expired or surplus Vidalista 80mg Yellow tablets should not be discarded by flushing. Investigate pharmaceutical take-back programs in your locality if possible. Absent such programs, render the tablets unusable (e.g., by mixing with undesirable materials like coffee grounds), seal them securely in a plastic bag, and then dispose of them in the regular household trash stream. **Lab Test Interference** The most likely effect on laboratory assessments relates to blood pressure measurements, as tadalafil can cause temporary reductions. It's prudent to inform medical personnel performing such tests that you are taking this potent 80mg formulation. Beyond this, significant interference with common blood or urine tests is generally not expected from tadalafil itself. **Handling Precautions** Use dry hands to handle the tablet and swallow it whole with water promptly after removing it from the packaging. Given the substantial 80 mg dosage, meticulous care must be taken to store this medication securely, well out of the reach and sight of children, pets, and any individuals (especially women) for whom it is not prescribed. **Counterfeit Identification Tips** **Packaging Authentication:** Thoroughly examine the external box and internal blister packs for signs of legitimacy: professional printing quality, accurate manufacturer details, clearly legible batch numbers and expiration dates, and intact tamper-evident features. Any discrepancies warrant caution. **Tablet Color and Consistency:** Genuine Vidalista 80mg Yellow tablets should exhibit a uniform yellow color and smooth coating. Be wary of variations in shade, presence of spots, cracks, crumbling texture, or poorly defined markings, as these could indicate a counterfeit product. **Source Legitimacy:** High-strength generic ED medications are frequent targets for counterfeiting. Procuring Vidalista 80mg Yellow from unverified online platforms or without a prescription significantly elevates the risk. Always use licensed pharmacies and validate the source. **Additional Advice** The distinctive yellow color is intended partly as a safety feature to help differentiate this high dose from other medications; always double-check you have the correct tablet. If traveling, carrying the medication in its original labeled packaging, potentially along with a prescription copy, is recommended to avoid complications. Always verify the expiration date before taking a tablet; efficacy and safety may be compromised past this date. Other formulations are available within the Vidalista product line to suit different patient requirements, including: Vidalista Black Vidalista Professional Vidalista CT Super Vidalista

---

## References

Drugs.com: Tadalafil. Drugs.com; 2024. [Updated Apr 22, 2024]. [Accessed May 21, 2024]. <https://www.drugs.com/tadalafil.html> MedlinePlus: Tadalafil. U.S. National Library of Medicine; 2024. [Updated Apr 15, 2024]. [Accessed May 21, 2024]. <https://medlineplus.gov/druginfo/meds/a604008.html> Mayo Clinic: Tadalafil (Oral Route) Description. Mayo Foundation for Medical Education and Research; 2024. [Updated Feb 1, 2024]. [Accessed May 21, 2024]. <https://www.mayoclinic.org/drugs-supplements/tadalafil-oral-route/description/drg-20067204> WebMD: Tadalafil Oral: Uses, Side Effects, Interactions, Pictures, Warnings & Dosing. WebMD; 2024. [Accessed May 21, 2024]. <https://www.webmd.com/drugs/2/drug-77875/tadalafil-oral/details>

---

## Disclaimer

Disclaimer: The information on this site is provided for informational purposes only and is not medical advice. It does not replace professional medical consultation, diagnosis, or treatment. Do not self-medicate based on the information presented on this site. Always consult with a doctor or other qualified healthcare professional before making any decisions about your health.