

iMedix: Your Personal Health Advisor.

Vidalista Professional

Vidalista Professional offers tadalafil in a rapidly dissolving tablet intended for quicker absorption, potentially leading to a faster onset of action for treating erectile difficulties. This formulation provides a convenient alternative for men seeking prompt readiness for sexual activity.

- **ActiveIngredient:** Tadalafil
 - **DosageForm:** Sublingual / Fast-Dissolving Tablet
 - **Dosage:** 20 mg
 - **Indications:** Employed in managing erectile dysfunction (ED)
 - **Manufacturer:** Varies (often Centurion Laboratories, verify packaging)
 - **Storage:** Keep below 30°C (86°F) in a moisture-free, dark environment. Retain in original packaging until use. Ensure it's kept away from children.
 - **Market Price:** 1.19
 - **Drug Status:** Prescription Only
-

Description

What is Vidalista Professional? Vidalista Professional is a medication utilized for addressing difficulties in achieving or maintaining an erection (erectile dysfunction) in adult men. It incorporates Tadalafil (20 mg) as its active pharmaceutical ingredient and is distinguished by its specialized fast-dissolving tablet format, designed for quicker entry into the bloodstream. Mechanism of Action The therapeutic effect of tadalafil stems from its role as a phosphodiesterase type 5 (PDE5) inhibitor. By interfering with the PDE5 enzyme's activity primarily in the penile tissues, tadalafil helps sustain higher levels of cyclic guanosine monophosphate (cGMP). This chemical messenger promotes the relaxation of arterial smooth muscles in the penis, allowing for increased blood influx when sexual stimulation occurs, thereby aiding erection development. The beneficial effects on erectile function can potentially span up to 36 hours following administration. Brand vs. Generic Status This product represents a branded generic version of tadalafil. While it relies on the well-established efficacy of tadalafil, the same active substance found within the reference product Cialis®, Vidalista Professional is distributed under its own unique trade name and formulation type, differing from the original brand. Available Forms Vidalista Professional is provided as sublingual or fast-dissolving tablets, each containing 20 milligrams of tadalafil. This dosage form is engineered to dissolve quickly when placed under the tongue, allowing the medication to be absorbed directly into the bloodstream, bypassing initial digestion in the stomach. Unique Features Rapid-Dissolve Formulation: The key characteristic is its fast-dissolving nature, enabling absorption through the tissues under the tongue. Potentially Quicker Onset: Due to sublingual absorption, the medication may begin to exert its effects more rapidly (often cited as within 15-30 minutes) compared to standard tablets that require digestion. Convenience and Discretion: The need to swallow with water may be bypassed, offering convenience and a more discreet method of administration. Standard Effective Dose: Contains 20 mg of tadalafil, a commonly prescribed and effective dose for many men experiencing ED.

General Instructions

How to Take Vidalista Professional Administration: Place one tablet under your tongue (sublingually). Allow the tablet to dissolve completely without chewing or swallowing it whole. Absorption occurs directly through the tissues under the tongue. Water is generally not needed to take the tablet, but you can drink water after it has fully dissolved if desired. **Food Considerations:** Vidalista Professional can generally be taken without regard to meals. Its primary absorption route bypasses the stomach. **Timing Guidelines Optimal Timing:** Administer the tablet approximately 15 to 30 minutes before you plan to engage in sexual activity, taking advantage of its rapid-dissolving design. Remember, sexual stimulation is required for it to work. **Duration of Effect:** The therapeutic benefits enabling erection can last for up to 36 hours after the tablet has dissolved and been absorbed. **Frequency Maximum Dosage:** Limit intake to no more than one Vidalista Professional tablet (20 mg) within any single 24-hour timeframe. **Missed Dose Situational Use:** This medication is intended for use as needed prior to sexual relations, not on a fixed daily regimen. If you anticipate sexual activity and forget to take it at the planned time, take it when you remember, as long as you adhere to the once-per-24-hours limit. There's no concept of "catching up" on a missed dose. **Storage Guidelines Conditions:** Store the tablets in a controlled room temperature setting, typically between 15°C and 30°C (59°F and 86°F). **Protection:** It is crucial to keep the tablets in their original blister packaging until the moment of use to safeguard them from air and moisture, which can degrade fast-dissolving formulations. Securely store away from children's reach and sight.

Side Effects

Using Vidalista Professional may lead to side effects in some individuals, similar to other medications containing tadalafil. While many people experience no issues or only minor ones, it's helpful to be aware of potential reactions. **Common Side Effects:** These reactions are encountered more frequently but are typically not severe and may diminish over time. **Headache:** A prevalent effect, often manageable with adequate hydration. **Facial Redness (Flushing):** A transient feeling of warmth spreading across the face. **Nasal Stuffiness:** Difficulty breathing through the nose or a sensation of congestion. **Digestive Discomfort:** Symptoms like indigestion or mild stomach upset may occur. **Musculoskeletal Pain:** Aches in the back or muscles can sometimes develop, usually appearing 12-24 hours post-dose and resolving within a couple of days. Simple pain relievers may help if cleared by your doctor. **Less Common Side Effects:** These are observed less often but warrant attention if they arise. **Lightheadedness:** A sense of dizziness or feeling off-balance. Refrain from operating vehicles or machinery if affected. **Visual Alterations:** Temporary changes such as slightly blurred vision or a subtle difference in color perception. **Rapid Pulse:** Noticing your heart beating quicker than its usual rhythm. **Ocular Irritation:** Discomfort or redness occurring in the eyes. **Serious Side Effects:** These situations are rare but demand prompt medical assistance. Cease using Vidalista Professional and contact emergency services or your doctor without delay if you notice: **Prolonged Erection (Priapism):** An erection that is painful or lasts continuously for over 4 hours. Urgent treatment is vital to prevent long-term penile injury. **Abrupt Vision Deterioration:** Sudden reduction or complete loss of sight affecting one or potentially both eyes (this could indicate NAION). **Sudden Hearing Impairment:** A rapid decrease in hearing ability, possibly accompanied by tinnitus (ringing ears) or vertigo. **Cardiac Symptoms During Intimacy:** Experiencing chest discomfort, significant dizziness, or feeling nauseous while engaging in sexual activity could indicate a serious heart issue, particularly for those with known cardiac conditions. **Severe Hypersensitivity Reaction:** Signs including skin eruptions (rash, hives), intense itching, trouble breathing, or significant swelling affecting the face, tongue, lips, or throat. **Management Note:** Should any side effect feel particularly troublesome, persist longer than expected, or cause worry, it is best to consult your healthcare professional for personalized advice. Avoid altering your medication regimen independently.

Uses

Approved Medical Application Vidalista Professional is designated for addressing erectile dysfunction (ED), a condition characterized by the persistent difficulty in attaining or sustaining an erection adequate for satisfactory sexual intercourse in adult men. **How It Addresses This Condition Facilitating Blood Inflow:** By

inhibiting the phosphodiesterase type 5 (PDE5) enzyme, tadalafil (the active component) helps preserve cGMP levels within the penile tissues. This enhances the natural process triggered by sexual arousal, where smooth muscles in penile arteries relax, permitting greater blood entry needed for an erection. **Rapid Absorption Design:** The sublingual (under the tongue) tablet formulation of Vidalista Professional is engineered for fast disintegration and absorption directly into the bloodstream. This characteristic may allow the medication to begin working more quickly than conventional tablets that must pass through the digestive system first. **Effective Dosage Strength:** The 20 mg tadalafil content represents a widely recognized and frequently prescribed dosage found effective for a broad range of men dealing with ED. **Extended Opportunity Window:** Consistent with tadalafil's properties, Vidalista Professional provides a lengthy timeframe, potentially up to 36 hours post-administration, during which achieving an erection upon stimulation is possible. **Important Considerations Symptom Management:** It's crucial to understand that Vidalista Professional helps manage the physical manifestation of ED but does not eliminate the underlying factors, which could be vascular, neurological, hormonal, or psychological in nature. **Arousal is Necessary:** The medication assists in the physiological response to sexual cues; it does not induce an erection spontaneously without concurrent sexual stimulation. **Formulation Advantage:** The fast-dissolving nature offers convenience, potentially faster onset, and an alternative for individuals who may have difficulty swallowing pills. **Professional Guidance:** Self-treatment is not advised. Consultation with a healthcare provider is essential to determine if Vidalista Professional is a suitable and safe option based on individual health status and other medications being taken. **Complementary Strategies:** For optimal management of ED, combining medication like Vidalista Professional with lifestyle modifications (e.g., exercise, healthy diet, stress reduction) and addressing any contributing health conditions is often recommended.

Safety advice

Interactions Alcohol:

- Use with caution
- Combining alcohol with Vidalista Professional might intensify side effects like feeling dizzy, experiencing headaches, or noticing a faster heartbeat, and could potentially lead to a drop in blood pressure. It is wise to moderate alcohol consumption significantly when planning to use this medication.

Interactions Other Medications:

- Consult your doctor
- Tadalafil can interact negatively with various other drugs, most critically with nitrates (which are absolutely forbidden). Inform your physician about all other treatments you use, including alpha-blockers, strong enzyme inhibitors or inducers (like certain antifungals or seizure medications), and blood pressure drugs, to prevent harmful effects such as severe hypotension.

Special Groups Pregnancy:

- Unsafe
- Vidalista Professional is formulated exclusively for use by adult men experiencing erectile dysfunction and is not intended or approved for administration to women, especially during pregnancy.

Special Groups Breastfeeding:

- Unsafe
- This medication should not be used by women, including those who are nursing. Tadalafil is designed for male sexual health conditions, and its use is inappropriate for breastfeeding mothers.

Special Groups Elderly:

- Use with caution
- Senior individuals (generally age 65+) might metabolize tadalafil differently or be more susceptible to its blood pressure-lowering effects and other side effects. A healthcare provider's assessment is necessary to confirm if Vidalista Professional is appropriate and safe for older patients.

Special Groups Children:

- Unsafe
- Use of Vidalista Professional is restricted to adult males (18 years or older). It must never be given to individuals younger than 18, as its safety and efficacy have not been studied in the pediatric population.

Effects on Activities Driving:

- Use with caution
- Potential side effects such as dizziness or alterations in vision could impair the ability to drive safely. Gauge your individual response to Vidalista Professional before getting behind the wheel, particularly upon initiating therapy.

Effects on Activities Operating Machinery:

- Use with caution
- Similar to driving, tasks requiring focus and coordination, like operating machinery, should be approached cautiously until you know how Vidalista Professional affects you, due to the possibility of dizziness or visual side effects.

Concerns

Key Concerns When Considering Vidalista Professional Cardiovascular Impact: Tadalafil, the active ingredient in Vidalista Professional, possesses blood pressure-lowering properties. Individuals with certain heart conditions, a recent history of stroke or heart attack, or those taking other blood pressure medications should use this drug cautiously, as there's a potential for symptomatic hypotension (dizziness, fainting). A thorough cardiovascular assessment by a doctor is recommended before use.

Potential for Priapism: Although uncommon, tadalafil use carries a risk of inducing priapism – an erection that persists painfully for more than four hours. This condition constitutes a medical emergency requiring immediate attention to avert potential long-term damage to erectile tissues.

Sensory Disturbances: Rare instances of sudden changes in vision (including a condition known as NAION) or sudden hearing loss have been associated with PDE5 inhibitors like tadalafil. Any abrupt alteration in sight or hearing should prompt discontinuation of the medication and urgent medical consultation.

Influence of Kidney or Liver Health: Compromised kidney or liver function can slow the body's elimination of tadalafil. This may lead to increased drug levels and a higher susceptibility to side effects. Dose adjustments or specific precautions might be necessary for individuals with renal or hepatic impairment, as directed by a healthcare provider.

When Not to Use Vidalista Professional Absolute Contraindications: This medication must not be used by individuals currently taking any form of nitrate medication (e.g., for heart conditions) due to the severe risk of dangerous hypotension. Other contraindications include a known allergy to tadalafil, specific severe cardiovascular conditions (like recent heart attack/stroke, unstable angina, significant heart failure), history of NAION-related vision loss, or severe liver problems. Use in patients with severe kidney disease may also be contraindicated or require significant caution.

Critical Reminders Proper Administration is Key: Understand that Vidalista Professional is formulated as a sublingual tablet designed to dissolve under the tongue for faster absorption. Swallowing it whole like a conventional tablet may alter its effectiveness and onset time.

Verify Your Source: Lifestyle medications like ED treatments are often counterfeited. Ensure you obtain Vidalista Professional from a legitimate pharmacy or supplier with a prescription to guarantee you receive the correct medication and dose, avoiding potentially harmful fakes.

Does Not Prevent STIs: Remember that Vidalista Professional solely addresses erectile function; it offers no protection against sexually transmitted infections. Practice safe sex measures as appropriate.

Communicate with Your Doctor: If side effects are persistent or particularly bothersome, or if the medication doesn't provide the desired results, discuss these issues with your healthcare provider. They can assess the situation and advise on potential alternatives or adjustments.

Warnings

Critical Health Authority Precautions Health regulators emphasize several crucial safety points regarding tadalafil, the active substance in Vidalista Professional: **Nitrate Co-administration Hazard:** A paramount warning highlights the potential for a drastic and unsafe lowering of blood pressure when tadalafil is used alongside any organic nitrate drug. **Assessment for Heart Conditions:** Due to its cardiovascular effects, a careful evaluation of a patient's heart health is deemed essential before initiating tadalafil therapy. **Risk of Acute Vision Loss (NAION):** Although infrequent, cases of non-arteritic anterior ischemic optic neuropathy (NAION), leading to sudden vision impairment, have been documented with this class of medication (PDE5 inhibitors). **Absolute Contraindications** Vidalista Professional is strictly unsuitable and should never be utilized under these conditions: **Use with Nitrates:** Simultaneous administration with any medication containing organic nitrates (like nitroglycerin sprays, patches, or tablets) or recreational "poppers" is absolutely forbidden due to life-threatening hypotensive risk. **History of Allergy:** Individuals who have previously experienced a significant allergic reaction (hypersensitivity) to tadalafil or any inactive ingredient within the Vidalista Professional tablet must avoid it. **Specific Severe Heart Problems:** Usage is contraindicated in patients diagnosed with significant cardiovascular instability, including recent (within months) myocardial infarction or stroke, ongoing unstable angina, certain classifications of heart failure, uncontrolled dangerous heart rhythms, or significantly low or high resting blood pressure. **Urgent Precautions** Should any of the following situations arise after taking Vidalista Professional, consider it a potential emergency requiring swift medical evaluation: **Priapism Occurrence:** An erection that persists beyond four hours without receding. This necessitates immediate medical care to prevent potentially permanent harm to penile erectile tissue. **Sudden Loss of Vision or Hearing:** Any abrupt change, decrease, or complete loss of sight in one or both eyes, or a sudden difficulty in hearing (possibly with tinnitus or dizziness). Stop the drug and seek help right away. **Cardiac Distress Symptoms:** Developing chest pain, acute dizziness, fainting sensations, or persistent nausea during or shortly after sexual relations. These could signal a serious cardiovascular response needing urgent assessment.

Dosage

Standard Dosing Guidelines Recommended Dose: Typically, one Vidalista Professional tablet, containing 20 mg of tadalafil, is taken as needed before sexual activity. It should be placed under the tongue to dissolve. **Maximum Frequency:** Adherence to a maximum limit is crucial: do not administer more than a single 20 mg tablet within a consecutive 24-hour interval. **Dose Adjustments Renal Impairment:** For mild to moderate kidney dysfunction, dosage adjustments for tadalafil 20 mg are not usually required when used 'as needed'. However, its use may need careful consideration or avoidance in cases of severe kidney disease (e.g., creatinine clearance below 30 mL/min) due to reduced drug clearance. Always follow physician recommendations. **Hepatic Impairment:** Patients with mild or moderate liver problems should exercise caution, and the 20 mg dose might be considered the maximum recommended 'as needed' dose under medical supervision. Vidalista Professional is generally not recommended for individuals with severe hepatic impairment (Child-Pugh Class C). **Elderly Patients:** While specific dose adjustments based solely on age are often unnecessary, older individuals might have underlying conditions (like reduced kidney function) or higher sensitivity to tadalafil's effects, warranting a careful review by a healthcare provider before using the 20 mg dose. **Missed Dose** As Vidalista Professional is intended for situational use before sexual activity, the concept of a 'missed dose' in a daily schedule does not apply. Simply use it when required, respecting the maximum 24-hour frequency guideline. **Overdose Management Symptoms:** Ingesting more Vidalista Professional than prescribed can intensify common side effects. Potential signs of overdose might include persistent headache, significant dizziness, fainting due to low blood pressure, prolonged erection (priapism), muscle aches, severe facial flushing, or substantial gastrointestinal upset. **Action:** In case of a suspected overdose, it is imperative to seek emergency medical assistance without delay or contact your local poison control center. Treatment will primarily involve supportive measures tailored to the patient's symptoms. **Critical Considerations Stick to Prescribed Amount:** Never independently decide to take more than one tablet thinking it will enhance the effect. Exceeding the 20 mg dose within 24 hours substantially elevates the risk

of adverse reactions. **Sublingual Absorption:** Recognize that the effectiveness and potentially faster onset are linked to the sublingual administration route. Ensure the tablet dissolves properly under the tongue for optimal results. **Communicate Effectiveness:** If the 20 mg dose seems inadequate or causes issues, consult your doctor. Do not attempt to compensate by taking additional tablets. Other doses or treatments might be more suitable.

Interactions

Drug-Drug Interactions Nitrates (Hazardous Combination): Taking Vidalista Professional concurrently with any form of nitrate medication (prescribed for chest pain, like nitroglycerin, or used recreationally as “poppers”) is absolutely contraindicated. This mix can cause a precipitous and dangerous decline in blood pressure. **Alpha-Blockers & Blood Pressure Agents:** When tadalafil is used alongside alpha-blockers (often for prostate issues or hypertension, e.g., terazosin) or other antihypertensive drugs, there’s an increased potential for additive blood pressure reduction. This may lead to symptoms like dizziness or fainting, particularly when standing up. Medical guidance on dose timing or adjustment might be necessary. **CYP3A4 Enzyme Modulators:** Medications that strongly inhibit the CYP3A4 metabolic pathway (e.g., certain antifungal drugs like itraconazole, HIV protease inhibitors like ritonavir) can substantially elevate tadalafil concentrations in the body, heightening side effect risks. Conversely, strong CYP3A4 inducers (like rifampin, certain anti-seizure drugs) can accelerate tadalafil’s elimination, potentially diminishing its effectiveness. Your doctor needs to know about these medications. **Other ED Treatments & Riociguat:** Combining Vidalista Professional with other phosphodiesterase type 5 inhibitors (such as sildenafil or vardenafil) or with riociguat (used for pulmonary hypertension) is not recommended and potentially unsafe due to the cumulative effects on blood pressure and increased risk of adverse events. **Drug-Food Interactions Heavy Meals:** While the sublingual absorption of Vidalista Professional largely bypasses the digestive system, potentially reducing the impact of food, extremely high-fat meals consumed around the same time might still theoretically slightly influence overall drug exposure or timing for some individuals, though likely less significantly than with standard oral tablets. **Grapefruit Products:** Grapefruit and its juice can interfere with the enzymes (specifically CYP3A4 in the intestinal wall) responsible for breaking down tadalafil. Consuming grapefruit could lead to higher levels of the drug in your system and an increased chance of side effects. Limiting or avoiding grapefruit is often suggested. **Alcohol:** Combining significant amounts of alcohol with Vidalista Professional is inadvisable. Alcohol can independently lower blood pressure and impair erectile function, and when taken with tadalafil, it can amplify dizziness, headache, rapid heartbeat, and the risk of postural hypotension (dizziness upon standing). **Drug-Supplement/Herb Interactions St. John’s Wort:** This herbal remedy can induce CYP3A4 enzymes, potentially lowering the blood levels and efficacy of tadalafil. Avoid using them together without medical advice. **Blood Pressure Affecting Supplements:** Supplements believed to influence blood pressure (e.g., L-arginine, high-dose fish oil) might theoretically have additive effects when taken with tadalafil; discuss supplement use with your healthcare provider. **Impact of Interactions** Interactions involving Vidalista Professional can manifest in various ways, ranging from diminished therapeutic outcomes (e.g., reduced drug levels) to amplified adverse effects (e.g., excessive blood pressure drop) or the emergence of serious complications requiring urgent care (especially with nitrates). **Critical Reminders Complete Medication History:** Providing your healthcare practitioner with an exhaustive list of all substances you ingest—including prescriptions, over-the-counter items, supplements, and recreational drugs—is crucial for effective interaction screening. **Counterfeit Interaction Dangers:** Be aware that counterfeit versions of Vidalista Professional pose a specific risk of unexpected drug interactions because they may contain undeclared or incorrect active ingredients, potentially reacting dangerously with your other medications. Always use legitimate suppliers. **Report Suspected Interactions:** If you experience any unusual or severe symptoms after taking Vidalista Professional, especially when combined with other substances, promptly inform your doctor, as this could indicate an adverse interaction requiring medical assessment.

FAQs

- **How does Vidalista Professional compare to Viagra in strength?**

Vidalista Professional (tadalafil) and Viagra (sildenafil) are both used to treat erectile dysfunction and work in similar ways, but they are not measured by "strength" in the traditional sense. While Viagra typically remains effective for about 4 to 6 hours, Vidalista Professional is designed to last much longer—up to 36 hours. This extended duration doesn't mean it's "stronger," but rather that it provides a longer window for spontaneity. The choice between them depends on individual preferences, health factors, and how your body responds.

- **Combining Vidalista Professional with recreational drugs: What are the risks?**

Mixing Vidalista Professional with recreational drugs (e.g., cocaine, poppers, or alcohol) can be dangerous. Tadalafil interacts with substances like nitrates or stimulants, potentially causing a sudden drop in blood pressure, fainting, or heart complications.

- **What happens if I double the dose of Vidalista Professional 20mg?**

Never take more than one 20mg tablet of Vidalista Professional in 24 hours. Doubling the dose increases the risk of severe side effects, including prolonged erections (priapism), dangerously low blood pressure, headaches, or vision changes. Stick to your prescribed dosage unless your doctor advises otherwise.

- **Is caffeine safe to consume with Vidalista Professional?**

Moderate caffeine intake (e.g., coffee or tea) is generally safe with tadalafil. However, excessive caffeine might worsen side effects like increased heart rate or jitteriness.

Other Details

Appearance Vidalista Professional 20 mg tablets are typically small, designed for rapid dissolution under the tongue. Their exact color (often white or off-white, but may vary) and markings (if any) depend on the specific manufacturer (frequently Centurion Laboratories, but verification is needed). Refer to the product packaging for precise identification details. Storage After Opening Retain each Vidalista Professional tablet in its original blister cavity until the moment of use. This is vital for protecting the fast-dissolving formulation from ambient moisture and air, which can compromise its integrity and dissolution properties. Do not store loose tablets or transfer them to pill organizers. Disposal Avoid disposing of unused or expired Vidalista Professional tablets by flushing them. Utilize community pharmaceutical take-back initiatives if accessible. If not, render the tablets unusable (e.g., mix with coffee grounds or cat litter), seal them in a bag, and discard them with regular household refuse to prevent accidental ingestion. Lab Test Interference Similar to other tadalafil products, Vidalista Professional can lower blood pressure, which might be reflected in readings taken during medical checks. Inform healthcare personnel conducting tests that you use this medication. Beyond potential blood pressure effects, significant interference with standard laboratory tests is generally not anticipated. Handling Precautions Use dry hands when handling the tablet. Place it directly under the tongue upon removal from the blister pack; do not chew or swallow it whole. Allow it to dissolve completely. Due to the active ingredient, keep this medication securely stored away from access by children, pets, or women. Counterfeit Identification Tips Packaging Scrutiny: Carefully inspect the box and blister packs for signs of authenticity – clear manufacturer information, legible batch codes and expiry dates, intact security seals or features. Substandard printing or damaged packaging could suggest a fake. Tablet Characteristics: Genuine fast-dissolving tablets should dissolve relatively quickly and smoothly under the tongue. Be suspicious of tablets that are unusually hard, gritty, dissolve very slowly, have an unexpected taste (unless flavored), or possess inconsistent appearance (color, shape, markings). Source Reliability: The risk of encountering counterfeit medication is significantly higher when purchasing from unverified online platforms or sources lacking proper accreditation. Obtain Vidalista Professional through legitimate channels requiring a prescription. Additional Advice The sublingual format may offer convenience for those who have difficulty swallowing conventional pills. If traveling, keeping the medication in its original marked packaging along with a prescription copy is advisable. Report any concerns about the tablet's quality (e.g., difficulty dissolving, unusual texture) to your pharmacist. Other formulations are available within the

Vidalista product line to suit different patient requirements, including: Vidalista Black Vidalista CT Super Vidalista

References

Drugs.com: Tadalafil. Drugs.com; 2024. [Updated Apr 22, 2024]. [Accessed May 21, 2024]. <https://www.drugs.com/tadalafil.html> MedlinePlus: Tadalafil. U.S. National Library of Medicine; 2024. [Updated Apr 15, 2024]. [Accessed May 21, 2024]. <https://medlineplus.gov/druginfo/meds/a604008.html> Mayo Clinic: Tadalafil (Oral Route) Description. Mayo Foundation for Medical Education and Research; 2024. [Updated Feb 1, 2024]. [Accessed May 21, 2024]. <https://www.mayoclinic.org/drugs-supplements/tadalafil-oral-route/description/drg-20067204> WebMD: Tadalafil Oral: Uses, Side Effects, Interactions, Pictures, Warnings & Dosing. WebMD; 2024. [Accessed May 21, 2024]. <https://www.webmd.com/drugs/2/drug-77875/tadalafil-oral/details>

Disclaimer

Disclaimer: The information on this site is provided for informational purposes only and is not medical advice. It does not replace professional medical consultation, diagnosis, or treatment. Do not self-medicate based on the information presented on this site. Always consult with a doctor or other qualified healthcare professional before making any decisions about your health.