

iMedix: Your Personal Health Advisor.

Vusion Ointment

Search for medical details concerning Vusion topical on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

Description

Side Effects Remember that this medication has been prescribed because the doctor has judged that the benefit to your child is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: increased irritation at the application site. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. How long does it take for this medicine to take effect? The onset of action of Vusion Ointment can vary depending on the severity of the diaper rash and individual response. Generally, noticeable improvement in symptoms such as redness and irritation may be observed within a few days of starting treatment. How long do the effects of this medicine last? The effects of Vusion Ointment typically last as long as the medication remains on the skin and continues to provide a protective barrier against moisture and irritants. Regular application as prescribed by a healthcare provider is necessary to maintain its therapeutic effects. Is it safe to consume alcohol while taking this medicine? Since Vusion Ointment is applied topically to the skin and not ingested, there are no known interactions with alcohol consumption. Therefore, it is generally safe to consume alcohol while using this medication. Is this a habit forming medicine? Vusion Ointment is not habit-forming as it does not contain any ingredients with addictive properties. It is used for short-term treatment of diaper rash and is not associated with dependence or abuse. Can this medicine be taken during pregnancy? The safety of Vusion Ointment during pregnancy has not been established. Pregnant individuals should consult with their healthcare provider before using this medication to assess the potential risks and benefits. Can this medicine be taken while breast-feeding? It is generally considered safe to use Vusion Ointment while breastfeeding, as the amount of active ingredients absorbed through the skin is minimal. However, it is recommended to avoid applying the ointment to areas of the breast that may come into contact with the infant's mouth to prevent accidental ingestion. It is advisable to consult with a healthcare provider before using this medication while breastfeeding to ensure safety for both the mother and the infant. **Uses** This product is used by children to treat diaper rash with a yeast infection, along with proper cleaning of the diaper area and frequent diaper changes. Miconazole is an azole

antifungal that works by preventing the growth of fungus. Zinc oxide and petrolatum work by forming a barrier on the skin to protect it from moisture and irritation. This product should not be used long-term or to prevent diaper rash. Unnecessary use or misuse of this product can lead to its decreased effectiveness. How to use Vusion Ointment Use this product on the skin only. Wash your hands before and after applying this product. Gently clean the diaper area with warm water and then pat dry. Gently apply a thin layer of this medication to the affected area as directed by your doctor, usually during each diaper change. Do not rub the medication into the skin or it may cause more irritation. This medication is usually used for 7 days. Do not apply the medication in the eyes, nose, mouth, or inside the vagina. If you do get the medication in those areas, flush with plenty of water. Use this medication regularly for the most benefit. Remember to use it with every diaper change. Continue to use this product for the full length of treatment, even if symptoms disappear after a few days. Stopping the medication too early may result in a return of the infection. Do not use this product for longer than prescribed. Tell the doctor if your child's condition lasts after 7 days of treatment or if it gets worse. Precautions Before using this product, tell the doctor or pharmacist if your child is allergic to miconazole, zinc oxide, or petrolatum; or to other azole antifungals (such as clotrimazole, ketoconazole); or if your child has any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell the doctor or pharmacist your child's medical history. During pregnancy, this medication should be used only when clearly needed. Pregnant women should use caution when applying this medication on their children. Discuss the risks and benefits with your doctor. It is unknown if this drug passes into breast milk. Breast-feeding women should use caution when applying this medication on their children. Consult your doctor before breast-feeding. Overdose This medicine may be harmful if swallowed. If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Do not share this medication with others. This medication should be used for your baby's current condition only. Do not use it later for another condition unless told to do so by the doctor. A different medication may be necessary in those cases. This product is not meant to replace frequent diaper changes. To help the diaper rash heal and to prevent future diaper rashes, check your baby's diaper often, and change the diaper whenever it appears wet/dirty. If you miss a dose, use it as soon as you remember. If it is near the time of the next diaper change, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products your child uses (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Side Effects

Uses

Interactions

Other Details