

iMedix: Your Personal Health Advisor.

Vutrisiran 25 Mg/0.5 ML Subcutaneous Syringe

Search for medical details concerning vutrisiran subcutaneous on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Bruising, redness, itching, or pain at the injection site may occur. Pain in the arm/leg, joint pain or shortness of breath may also occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. This medication decreases vitamin A levels in your blood. Ask your doctor about taking a vitamin A supplement. Tell your doctor right away if you develop signs of low vitamin A levels, especially vision changes such as dry eyes, blurred vision, or trouble seeing in dim light (night blindness). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses Vutrisiran is used to treat nerve problems due to a certain inherited condition (transthyretin-mediated amyloidosis). This condition causes a protein that your body normally makes (transthyretin-TTR), to change shape and build up in different parts of your body, which can affect the nerves, heart, and gut. This medication works by decreasing the buildup of TTR in your body. How to use Vutrisiran 25 Mg/0.5 ML Subcutaneous Syringe This medication is given by injection under the skin by a health care professional. It is injected in the thigh, abdomen, or upper arm as directed by your doctor, usually once every 3 months. Use this medication regularly to get the most benefit from it. To help you remember, mark the days on the calendar when you need to receive this medication. Tell your doctor if your condition does not get better or if it gets worse. Precautions Before using vutrisiran, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Discuss the risks and benefits with your doctor. It is unknown if this medication passes into breast milk. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or

trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Lab and/or medical tests may be done while you are using this medication. Keep all medical and lab appointments. It is important to get each dose of this medication as scheduled. If you miss a dose, ask your doctor or pharmacist right away for a new dosing schedule. Not applicable. This medication is given in a hospital or clinic or doctor's office and will not be stored at home. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval.

Side Effects

Uses

Interactions

Other Details
