

# iMedix: Your Personal Health Advisor.

## Wilzin 25 Mg (Zinc) Capsule Wilson's Disease Treatment Agents

---

Search for medical details concerning Wilzin oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
- **DosageForm:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

### Description

Side Effects Upset stomach may occur. If this effect lasts or gets worse, tell your doctor or pharmacist promptly. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. How long does it take for this medicine to take effect? The onset of action of Wilzin 25 mg capsules, containing the active ingredient zinc, can vary depending on the condition being treated and individual factors. Generally, it may take several days to weeks of regular use before the therapeutic effects become noticeable. How long do the effects of this medicine last? The duration of effects of Wilzin 25 mg capsules also varies depending on factors such as the dosage, frequency of administration, and the specific condition being treated. Generally, the effects may persist for a period of time after discontinuation of the medication, especially if used for addressing zinc deficiency. Is it safe to consume alcohol while taking this medicine? There are no known interactions between alcohol consumption and Wilzin capsules. However, it is advisable to consult with a healthcare professional regarding potential interactions, as excessive alcohol intake may interfere with zinc absorption and metabolism. Is this a habit forming medicine? Wilzin 25 mg capsules are not known to be habit-forming. They are typically used for addressing zinc deficiency or as a dietary supplement and are not associated with the development of physical dependence or addiction. Can this medicine be taken during pregnancy? Zinc is an essential nutrient for pregnant women, and maintaining adequate zinc levels is important for maternal and fetal health. Wilzin capsules may be prescribed during pregnancy if there is a diagnosed zinc deficiency, but it is essential to consult with a healthcare provider before initiating any medication during pregnancy. Can this medicine be taken while breast-feeding? Zinc is also important for lactating women to support milk production and the health of both the mother and the infant. Wilzin capsules may be considered safe to use while breastfeeding, but it is important to consult with a healthcare provider to assess the individual's zinc status and determine the appropriate dosage and duration of use during breastfeeding. Uses This medication is used after other

medications to treat a certain liver disease (Wilson's disease). This inherited disease causes the liver to hold onto too much copper, resulting in liver damage and other serious problems. This medication causes the intestines to make more of a certain substance (a protein) that prevents the body from absorbing too much copper from food, thereby preventing further damage. How to use Wilzin 25 Mg (Zinc) Capsule Wilson's Disease Treatment Agents Take this medication by mouth as directed by your doctor, usually 3 times daily. Take each dose at least 1 hour before or 2 hours after all food or beverages (except water). Swallow the capsule whole. Do not crush or chew the capsules. Dosage is based on your medical condition and response to treatment. Use this medication regularly to get the most benefit from it. To help you remember, take it at the same times each day. Tell your doctor if your condition does not improve or if it worsens. Precautions Before taking zinc acetate, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Tell your doctor if you are pregnant before using this medication. This drug passes into breast milk and may have undesirable effects on a nursing infant. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include: nausea, vomiting, stomach/abdominal pain, mental changes, decrease in urine production. Do not share this medication with others. Lab and/or medical tests (such as 24-hour urine copper level, liver function, speech evaluation) should be done while you are taking this medication. Keep all medical and lab appointments. Consult your doctor for more details. If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Take your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

**Interactions** Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. This product can decrease the absorption of other drugs such as tetracycline antibiotics (such as doxycycline, minocycline), bisphosphonates (such as alendronate), and quinolone antibiotics (such as ciprofloxacin, levofloxacin). Separate your doses of these medications as far as possible from your doses of this product. Ask your doctor or pharmacist about how long you should wait between doses and for help finding a dosing schedule that will work with all your medications.

---

---

## **Side Effects**

---

## **Uses**

---

---

---

---

---

## Interactions

---

---

---

---

## Other Details

---

---