

iMedix: Your Personal Health Advisor.

Zinc-15

Search for medical details concerning Zinc-15 oral on iMedix including its uses, side effects and safety, interactions, pictures, warnings and user ratings.

- **ActiveIngredient:**
 - **DosageForm:**
 - **Dosage:**
 - **Indications:**
 - **Manufacturer:**
 - **Storage:**
 - **Market Price:**
 - **Drug Status:**
-

Description

Side Effects Nausea, stomach upset, heartburn may occur. If any of these effects last or get worse, tell your doctor or pharmacist promptly. If your doctor has directed you to use this product, remember that your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this product do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: signs of infection (such as sore throat that doesn't go away, fever, chills), sores in mouth, unusual tiredness, weakness. A very serious allergic reaction to this product is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at www.fda.gov/medwatch. In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Uses This product is a mineral used to treat or prevent low levels of zinc. How to use Zinc-15 Take this product by mouth as directed by your doctor or the product package. Take this product 1 hour before or 2 hours after meals. It may be taken with food if it upsets your stomach. Avoid milk, bran, grains, or cereals within 2 hours of taking this product. Use this product regularly in order to get the most benefit from it. To help you remember, use it at the same time(s) each day. Precautions Before taking this product, tell your doctor or pharmacist if you have any allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this product, tell your doctor or pharmacist your medical history, especially of: low blood levels of copper. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). Tell your doctor if you are pregnant before using this product. This product passes into breast milk. While there have been no reports of harm to nursing infants, consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include severe nausea, vomiting, dizziness, fainting, and shortness of breath. It is important to maintain a well balanced diet while taking this medicine. Foods containing good sources of zinc include lean red meats, seafood, peas, and beans. Lab tests (such as zinc and copper levels) should be done while you are taking this product. Keep all

medical and lab appointments. Consult your doctor for more details. If you miss a dose, use it as soon as you remember. If it is near the time of the next dose, skip the missed dose. Use your next dose at the regular time. Do not double the dose to catch up. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. Some drugs that may interact with this product are: penicillamine, phosphorus containing products. This product can decrease the absorption of other drugs such as tetracycline antibiotics (including doxycycline, minocycline), bisphosphonates (such as alendronate), and quinolone antibiotics (such as ciprofloxacin, levofloxacin). Separate your doses of these medications as far as possible from your doses of this product. Ask your doctor or pharmacist about how long you should wait between doses and for help finding a dosing schedule that will work with all your medications.

Side Effects

Uses

Interactions

Other Details
