

# iMedix: Your Personal Health Advisor.

## Zolpidem

---

Zolpidem is a medication primarily used to treat short-term sleep disturbances, such as difficulty falling or staying asleep, which is often referred to as insomnia.

- **Active Ingredient:**
- **Dosage Form:**
- **Dosage:**
- **Indications:**
- **Manufacturer:**
- **Storage:**
- **Market Price:**
- **Drug Status:**

---

## Description

**Side Effects** Dizziness may occur. If this effect lasts or gets worse, tell your doctor or pharmacist promptly. This medication may make you sleepy during the day. Tell your doctor if you have daytime drowsiness. Your dose may need to be adjusted. Remember that this medication has been prescribed because your doctor has judged that the benefit to you is greater than the risk of side effects. Many people using this medication do not have serious side effects. Tell your doctor right away if you have any serious side effects, including: memory loss, mental/mood/behavior changes (such as new/worsening depression, abnormal thoughts, thoughts of suicide, hallucinations, confusion, agitation, aggressive behavior, anxiety). A very serious allergic reaction to this drug is rare. However, get medical help right away if you notice any symptoms of a serious allergic reaction, including: rash, itching/swelling (especially of the face/tongue/throat), severe dizziness, trouble breathing. This is not a complete list of possible side effects. If you notice other effects not listed above, contact your doctor or pharmacist. In the US – Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or at [www.fda.gov/medwatch](http://www.fda.gov/medwatch). In Canada – Call your doctor for medical advice about side effects. You may report side effects to Health Canada at 1-866-234-2345. Zolpidem is used for a short time to treat a certain sleep problem (insomnia) in adults. If you have trouble falling asleep, it helps you fall asleep faster, so you can get a better night's rest. Zolpidem belongs to a class of drugs called sedative-hypnotics. It acts on your brain to produce a calming effect. How to use zolpidem oral Read the Medication Guide and, if available, the Patient Information Leaflet provided by your pharmacist before you start taking zolpidem and each time you get a refill. If you have any questions, ask your doctor or pharmacist. Take this medication by mouth on an empty stomach as directed by your doctor, usually once a night. Since zolpidem works quickly, take it right before you get into bed. Do not take it with or after a meal because it will not work as quickly. If you are taking the capsule form of this medication, swallow whole. Do not open, chew, or crush the capsules. Do not take a dose of this drug unless you have time for a full night's sleep of at least 7 to 8 hours. If you have to wake up before that, you may have some memory loss and may have trouble safely doing any activity that requires alertness, such as driving or operating machinery. ( ) Dosage is based on your gender, age, medical condition, other medications you may be taking, and response to treatment. Do not increase your dose, take it more often, or use it for longer than prescribed. Do not take more than 10 milligrams a day. Women are usually prescribed a lower dose because the drug is removed from the body more slowly than in men. Older adults are usually prescribed a lower dose to decrease the risk of side effects. If you suddenly stop using this medication, you

may have withdrawal symptoms (such as nausea, vomiting, flushing, stomach cramps, nervousness, shakiness). To help prevent withdrawal, your doctor may lower your dose slowly. Withdrawal is more likely if you have used zolpidem for a long time or in high doses. Tell your doctor or pharmacist right away if you have withdrawal. Though it helps many people, this medication may sometimes cause addiction. This risk may be higher if you have a substance use disorder (such as overuse of or addiction to drugs/alcohol). Take this medication exactly as prescribed to lower the risk of addiction. Ask your doctor or pharmacist for more details. When this medication is used for a long time, it may not work as well. Talk with your doctor if this medication stops working well. Tell your doctor if your condition lasts after 7 to 10 days, or if it gets worse. You may have trouble sleeping the first few nights after you stop taking this medication. This is called rebound insomnia and is normal. It will usually go away after 1-2 nights. If this effect continues, contact your doctor. Warnings Rarely, after taking this drug, people have gotten out of bed and driven vehicles while not fully awake ("sleep-driving"). People have also sleepwalked, prepared/eaten food, made phone calls, or had sex while not fully awake. Often, these people do not remember these events. This problem can be dangerous (possibly fatal) to you or to others. If you find out that you have done any of these activities after taking this medication, tell your doctor right away. You should not take this medication or similar medications (such as eszopiclone, zaleplon) if you have this reaction to the medication. Interactions Drug interactions may change how your medications work or increase your risk for serious side effects. This document does not contain all possible drug interactions. Keep a list of all the products you use (including prescription/nonprescription drugs and herbal products) and share it with your doctor and pharmacist. Do not start, stop, or change the dosage of any medicines without your doctor's approval. A product that may interact with this drug is: sodium oxybate. Other medications can affect the removal of zolpidem from your body, which may affect how zolpidem works. Examples include azole antifungals (such as ketoconazole), rifampin, St. John's Wort, among others. The risk of serious side effects (such as slow/shallow breathing, severe drowsiness/dizziness) may be increased if this medication is taken with other products that may also cause drowsiness or breathing problems. Tell your doctor or pharmacist if you are taking other products such as opioid pain or cough relievers (such as codeine, hydrocodone), alcohol, marijuana (cannabis), other drugs for sleep or anxiety (such as alprazolam, lorazepam, zopiclone), muscle relaxants (such as carisoprodol, cyclobenzaprine), or antihistamines (such as cetirizine, diphenhydramine). Check the labels on all your medicines (such as allergy or cough-and-cold products) because they may contain ingredients that cause drowsiness. Ask your pharmacist about using those products safely. Precautions Before taking zolpidem, tell your doctor or pharmacist if you are allergic to it; or if you have any other allergies. This product may contain inactive ingredients, which can cause allergic reactions or other problems. Talk to your pharmacist for more details. Before using this medication, tell your doctor or pharmacist your medical history, especially of: kidney disease, liver disease, mental/mood problems (such as depression, thoughts of suicide), personal or family history of a substance use disorder (such as overuse of or addiction to drugs/alcohol), personal or family history of sleepwalking, lung/breathing problems (such as chronic obstructive pulmonary disease-COPD, sleep apnea), a certain muscle disease (myasthenia gravis). The effects of this drug can last even after you wake up the next day. If you did not get 7 to 8 hours of sleep or took other medications that made you sleepy or are more sensitive to this drug, you may feel alert but not think clearly enough to drive. You may also experience dizziness or blurred/double vision. Alcohol or marijuana (cannabis) can make you more dizzy. Wait at least 8 hours after taking this drug before driving, and do not drive, use machinery, or do anything that needs alertness until you can do it safely. This medication may also increase the risk of falls. Avoid alcoholic beverages. Talk to your doctor if you are using marijuana (cannabis). Children may be more sensitive to the side effects of this drug, especially dizziness and hallucinations. Older adults may be more sensitive to the side effects of this drug, especially dizziness, confusion, unsteadiness, and excessive drowsiness. These side effects can increase the risk of falling. Before having surgery, tell your doctor or dentist about all the products you use (including prescription drugs, nonprescription drugs, and herbal products). During pregnancy, this medication should be used only when clearly needed. Babies born to mothers who have used this drug late in the last 3 months of pregnancy may develop unusual sleepiness, trouble breathing, unusual limpness, or withdrawal symptoms. Get medical help right away if you notice any unusual symptoms in your newborn. Discuss the risks and benefits with your doctor. A small amount of this medication passes into breast milk and may have undesirable effects on a nursing infant (such as unusual sleepiness, trouble breathing, or unusual limpness). Get medical help right away if you notice any unusual

symptoms in your baby. Ask your doctor if you should pump and discard your breast milk during treatment and for 23 hours after a dose of this medication to lessen the risk of these effects in your baby. Consult your doctor before breast-feeding. Overdose If someone has overdosed and has serious symptoms such as passing out or trouble breathing, call 911. Otherwise, call a poison control center right away. US residents can call their local poison control center at 1-800-222-1222. Canada residents can call a provincial poison control center. Symptoms of overdose may include slowed breathing or a deep sleep from which you cannot be awakened. Do not share this medication with others. Sharing it is against the law. As you get older, your sleep pattern may naturally change and your sleep may be interrupted several times during the night. Consult your doctor or pharmacist for ways to improve your sleep without medication, such as avoiding caffeine and alcohol close to bedtime, avoiding daytime naps, and going to bed at the same time each night. If you miss a dose, do not take it unless you have time to sleep for 7 to 8 hours afterwards. Store at room temperature away from light and moisture. Do not store in the bathroom. Keep all medications away from children and pets. Do not flush medications down the toilet or pour them into a drain unless instructed to do so. Properly discard this product when it is expired or no longer needed. Consult your pharmacist or local waste disposal company.

---

---

---

## **Side Effects**

---

---

---

## **Uses**

---

---

---

---

---

---

---

## **Interactions**

---

---

---

---

## **Other Details**

---

---

---